

Geostatistics with compositional data, an overview

Tolosana-Delgado, R.; Mueller, U.; van den Boogaart, K. G.; Pawlowsky-Glahn, V.;
Egozcue, J. J.;

Originally published:

November 2018

Mathematical Geosciences 51(2019), 485-526

DOI: <https://doi.org/10.1007/s11004-018-9769-3>

Perma-Link to Publication Repository of HZDR:

<https://www.hzdr.de/publications/Publ-26318>

Release of the secondary publication
on the basis of the German Copyright Law § 38 Section 4.

Geostatistics with compositional data, an overview

R. Tolosana-Delgado¹, U. Mueller² and K.G. van den Boogaart^{1,3}

¹Helmholtz-Zentrum Dresden-Rossendorf,
Helmholtz Institute Freiberg for Resources Technology,
Freiberg, Germany

r.tolosana@hzdr.de, boogaart@hzdr.de

²School of Science,
Edith Cowan University,
Joondalup (WA), Australia, u.mueller@ecu.edu.au

³Institute for Stochastics,
Technical University “Bergakademie” Freiberg, Germany

December 15, 2017

Abstract

This paper presents an overview of results for the geostatistical analysis of collocated multivariate data sets, whose variables form a composition, i.e. the components represent the relative importance of the parts forming a whole. Such data sets occur most often in mining, hydro-geochemistry and soil science, but the results gathered here are relevant for any regionalised compositional data set. The paper covers the basic definitions, the analysis of the spatial codependence between components, mapping methods of cokriging and cosimulation honoring compositional constraints, the role of pre- and post-transformations like log-ratios or multivariate normal score transforms, and block-support upscaling. The main result is that log-ratio scores can and should be analysed jointly with multivariate geostatistics techniques, in which case the system data-variograms-cokriging enjoys an intrinsic consistency, delivering the same results regardless of which log-ratio transformation was used to represent them. Proofs of all statements are included in an appendix.

1 Introduction

Geostatistics is the name commonly given to a set of tools and concepts for modeling data exhibiting codependence related to the spatial proximity of their sampling locations. Two-point geostatistics was based on the concept of covariance/correlation function. A covariance (correlation) function describes the evolution of the covariance (correlation) exhibited by two variables geographically a lag distance h apart, as a function of multiples of this lag. Under reasonable assumptions (ergodicity, spatial stationarity, some sort of anisotropy and Gaussianity, eventually after a point-wise transformation), such a function can be estimated and modelled from geo-referenced data sets, and further used to obtain maps (interpolations or simulations) of the studied variables. Since the founding works of Matheron (1963, 1965, 1971), Geostatistics has become a standard method in many fields of geosciences (Journel and Huijbregts, 1978; Cressie, 1991; Chilés and Delfiner, 1999; Wackernagel, 2003).

A data set is compositional if it is formed by several variables which jointly describe the relative weight, importance or influence of a set of components forming a system. It is well-known in the geosciences (Chayes, 1960) that compositional data are affected by the spurious correlation problem, namely that the apparent correlation between two components of the system depends on what other components are considered in the system. As a consequence the common interpretation of correlation is no longer valid, that is, correlation does not represent a valid measure of linear association between two components. This carries over to geo-referenced compositions, as their spatial auto and cross-correlation functions are as spurious as correlation coefficients are for non-regionalized compositions (Pawłowsky, 1984).

Attending to certain consistency principles and basic mathematical tractability, Aitchison (1982) proposed to deal with compositional data after a log-ratio transformation. His methodology was first applied to regionalized compositions by Pawłowsky (1986). Several case studies were presented afterwards (Pawłowsky-Glahn and Burger, 1992; Pawłowsky, 1989; Pawłowsky-Glahn et al., 1995; Lark and Bishop, 2007; Morales Boezio, 2010; Ward and Mueller, 2012; Morales Boezio et al., 2012; van den Boogaart and Tolosana-Delgado, 2003; Ward and Mueller, 2013; Sun et al., 2014), however the geostatistical community has not yet accepted log-ratio methods, even after the appearance of a monograph (Pawłowsky-Glahn and Olea, 2004), further theoretical developments (Pawłowsky-Glahn, 2003; Tolosana-Delgado, 2006; Tolosana-Delgado and Boogaart, 2013) and at least one practical guide (Tolosana-Delgado et al., 2011).

Apart from practical reasons (lack of implementation of the proposed methods in commercial software), some theoretical concerns might also explain that reluctance: the possible effect of the particular log-ratio transformation chosen, the alleged bias of estimates obtained, the supposed lack of upscaling models and block kriging methods, and the widespread tradition of applying a gaussian anamorphosis (or normal scores transform). This contribution presents a comprehensive discussion of the fundamentals of log-ratio methods for geostatistical applications, attending to each one of these issues. An illustration case study is collated through the text, to illustrate the concepts, methods and tools together with their theoretical discussion.

Section 2 introduces this illustration case study. Section 3 reviews the grounding ideas, principles, concepts and operations of compositional data analysis relevant for geostatistical applications. Section 4 summarizes the common geostatistical practice and existing alternative methods for the spatial modeling compositional data, particularly focusing on their problems and caveats. Section 5 introduces several ways to specify the spatial dependence between the components of a regionalized composition, and the links between them. Section 6 presents some considerations on cokriging and cross-validation, while Section 6.2 discusses the usages of the conditional distribution for cosimulation and estimation of non-linear quantities, such as block cokriging for compositional data. Proofs of the relevant statements are referenced to previous works or included in the first appendix. A second appendix includes a concise, practical workflow.

2 Illustration: the K-Pit data set

The K-Pit is a high-grade iron ore deposit (Angerer and Hagemann, 2010) of Banded Iron Formation (BIF) type, in the Archean Koolyanobbing Greenstone Belt, Western Australia. The deposit is located 360km east of Perth in the Southern Cross Province of the Yilgarn Craton. The greenstone belt strikes northwest and is 35km long and 8 km wide, approximately. It is composed of folded sequences of amphibolites, meta-komatiites and intercalated metamorphosed BIF (Griffin, 1981). This particular deposit occurs where the main BIF horizon is offset by younger striking faults (Angerer and Hagemann, 2010). It consists of several mineralogical-textural types, including hard high-grade magnetite, hematite and goethite ores and medium-grade fault-controlled hematite-quartz breccia, as well as hematite-magnetite BIF.

Three domains, 202 (west main domain), 203 (east main domain) and 300 (hematite hanging wall), were selected as they can be considered reasonably

homogeneous from a mineralogical point of view: the iron-rich facies is dominated by hematite in all of them, with minor contributions of magnetite or goethite/limonite.

This study focuses on the geostatistical modelling of the percentages of Al_2O_3 , Fe, Mn, P and SiO in 2495 samples of support 2m long from 226 drill holes within the 3 zones. Their conventional statistics are shown in Table 1.

Table 1: Traditional statistics of the K-Pit data set

	Minimum	Maximum	Mean	Variance	Skew
Al_2O_3	0.023	25.000	0.836	2.672	7.347
Fe	18.040	69.521	61.979	20.521	-3.124
Mn	0.001	1.694	0.054	0.008	8.655
P	0.003	0.663	0.121	0.004	1.219
SiO_2	0.170	46.580	3.582	23.441	3.240
Sum	45.490	82.015	66.572	7.841	0.253

3 Compositional data analysis

3.1 The concepts of composition

A data set is compositional whenever its variables inform of the relative contribution of a set of components forming a whole. “*Relative importances*” cannot be negative numbers. It is often said that their total sum is bounded by a unit constant κ , such as 100%, 1 or 10^6 ppm, although this is not a necessary condition. If we denote by $\mathbf{z} = [z_1, z_2, \dots, z_D]$ a vector of D components, then

$$z_i \geq 0 \quad \text{and} \quad \sum_{i=1}^D z_i \leq \kappa. \quad (1)$$

The set \mathcal{S}^D of all points of D -dimensional real space \mathbb{R}^D satisfying these conditions is called the D -part *simplex*. The sum equality is fulfilled if all components of the whole are being considered, as the sum of contributions of an exhaustive partition must account for “everything”.

If the sum is considered informative and is always below κ , it is possible to complement the composition with a further $(D + 1)$ -th component, the fill-in variable

$$z_{D+1} = \kappa - \sum_{i=1}^D z_i. \quad (2)$$

This option is particularly interesting if the fill-in variable has a physical interpretation, as happens with the K-pit case study.

If the composition does not satisfy the constant sum constraint, but its total sum is deemed uninformative, then the constraint can be forced by applying the closure operation, denoted as $\mathcal{C}[\cdot]$ and defined as

$$\mathcal{C}[\mathbf{z}] = \frac{1}{\sum_{i=1}^D z_i} \mathbf{z}, \quad (3)$$

effectively removing the uninformative variability from the total sum. This indeed opens the door to a more general definition of composition: if two vectors \mathbf{x} and \mathbf{y} of positive components are such that $\mathcal{C}[\mathbf{x}] = \mathcal{C}[\mathbf{y}]$, they are said to be *compositionally equivalent*, and the class of all vectors \mathbf{y} compositionally equivalent to \mathbf{x} is called a *composition* (Barceló-Vidal, 2003). In fact, analyzing the original vector \mathbf{x} or its closed version $\mathcal{C}[\mathbf{x}]$ should give the same results, being compositionally equivalent. This case occurs, for instance, in the study of geochemical surveys with partial leachates or of vegetal matrices, because the total sum may then be related to sample preparation issues or of the age of the plants.

Sometimes, compositional data are expressed in non-dimensionless units (mg/L, mol/Kg, meq/L) which do not appear to satisfy either of the two cases. For instance, this is often the case in groundwater reservoir quality studies. In those cases it should be still possible to conceive a meaningful change to dimensionless units (mostly Kg/Kg or mol/mol) in which one could decide which of the two preceding cases apply. In many of these cases (van den Boogaart and Tolosana-Delgado, 2003), the theory of compositional analysis ensures that final results with dimensionless units or with the original units are mutually consistent.

When only a subset of components is considered, one speaks of a *subcomposition*. Typically, subcompositions are reclosed to recast their information relative to the total that the selected variables represent. Building/closing a subcomposition is routinely done in geology when representing 3 components of a composition in a ternary diagram.

To work with compositional data Aitchison (1986) proposed two principles, subcompositional coherence and scale invariance, that should be satisfied by any meaningful method of analysis in order to avoid the risks of spurious correlation. Subcompositional coherence implies that results obtained analysing a subcomposition cannot contradict those obtained analysing the whole composition. Scale invariance proposes that results should be the same for a composition \mathbf{z} and any scaled version $p\mathbf{z}$, such as the composition resulting from a change of

units from % to ppm. Both conditions imply working with ratios. This holds for all three cases of compositions mentioned: with informative total sum, with uninformative total sum, and with non-dimensionless units.

3.2 Geometry

The simplex \mathcal{S}^D , the sample space of compositions, can be given an inner product space structure (Pawlowsky-Glahn and Egozcue, 2001). The operations addition and scalar multiplication in the underlying vector space structure are given by the operations of perturbation (\oplus Aitchison, 1982) and powering (\odot Aitchison, 1982), defined respectively as

$$\begin{aligned}\mathbf{a} \oplus \mathbf{b} &= \mathcal{C}[a_1 b_1, a_2 b_2, \dots, a_D b_D], \\ \lambda \odot \mathbf{a} &= \mathcal{C}[a_1^\lambda, a_2^\lambda, \dots, a_D^\lambda].\end{aligned}\quad (4)$$

An inner product (Pawlowsky-Glahn and Egozcue, 2001) on this vector space is given by

$$\langle \mathbf{a}, \mathbf{b} \rangle_A = \frac{1}{D} \sum_{i>j}^D \log \frac{a_i}{a_j} \log \frac{b_i}{b_j}.\quad (5)$$

This induces a metric, called the Aitchison distance (Aitchison, 1986)

$$d_A(\mathbf{a}, \mathbf{b}) = \sqrt{\langle \mathbf{a} \ominus \mathbf{b}, \mathbf{a} \ominus \mathbf{b} \rangle_A} = \sqrt{\frac{1}{D} \sum_{i>j}^D \left(\log \frac{a_i}{a_j} - \log \frac{b_i}{b_j} \right)^2}.\quad (6)$$

In his seminal works about compositional data analysis, Aitchison (1982, 1986) defined the additive log-ratio transformation (alr), the centered log-ratio transformation (clr), and their inverses, respectively, as

$$\begin{aligned}\text{alr}(\mathbf{z}) &= \log \frac{\mathbf{z}_{-D}}{z_D} = \mathbf{J} \cdot \log \mathbf{z} = \mathbf{y}^a, & \text{alr}^{-1}(\mathbf{y}^a) &= \mathcal{C}[\exp[\mathbf{y}^a; 0]], \quad (7) \\ \text{clr}(\mathbf{z}) &= \log \frac{\mathbf{z}}{\sqrt[D]{z_1 z_2 \cdots z_D}} = \mathbf{H} \cdot \log \mathbf{z} = \mathbf{y}^c & \text{clr}^{-1}(\mathbf{y}^c) &= \mathcal{C}[\exp(\mathbf{y}^c)], \quad (8)\end{aligned}$$

where \mathbf{z}_{-D} is the original composition without the last component, $[\mathbf{y}^a; 0]$ is a vector expanded with a zero component, logarithms are applied component-wise, and

$$\begin{aligned}\mathbf{J} &= [\mathbf{I}_{(D-1) \times (D-1)}; -\mathbf{1}_{D-1}], \\ \mathbf{H} &= \mathbf{I}_{D \times D} - \frac{1}{D} \mathbf{1}_{D \times D}.\end{aligned}\quad (9)$$

In these and the following expressions, subscripts of a matrix or vector indicate their size, and \mathbf{I} is the identity matrix. Note that the row and column sums of \mathbf{H} are equal to 0, which makes the clr coefficients sum to zero too.

With the introduction of the inner product space structure, it was recognized that alr-transformed scores are nothing other but coordinates of compositions with respect to a non-orthogonal basis, and clr-scores are coefficients in a generating system of \mathcal{S}^D . In general terms, a composition and its coordinates in any basis are related by

$$\mathbf{y} = \mathbf{\Psi} \cdot \log \mathbf{z}, \quad \mathbf{z} = \mathcal{C} [\exp(\mathbf{\Psi}^- \cdot \mathbf{y})], \quad (10)$$

where $\mathbf{\Psi}$ is a $(D - 1) \times D$ -matrix whose rows sum to zero

$$\mathbf{\Psi} \cdot \mathbf{1}_D = \mathbf{0}_{D-1}, \quad (11)$$

and $\mathbf{\Psi}^-$ is its Moore-Penrose generalized inverse. Note that both matrices \mathbf{J} and \mathbf{H} satisfy these conditions. The columns of $\mathbf{\Psi}^-$ are the clr-transformed coefficients of the vectors of the orthonormal basis used.

Following this realization, Egozcue et al. (2003) proposed using orthonormal coordinate systems, to avoid having to work with generalized inverses. If the coordinate system is orthonormal, then matrix $\mathbf{\Psi}$ satisfies

$$\mathbf{\Psi} \cdot \mathbf{\Psi}^t = \mathbf{I}_{(D-1) \times (D-1)}, \quad \mathbf{\Psi}^t \cdot \mathbf{\Psi} = \mathbf{H}. \quad (12)$$

It follows that $\mathbf{\Psi}^- = \mathbf{\Psi}^t$, and the isometric log-ratio transformation (and its inverse) can be defined as

$$\mathbf{y} = \text{ilr}_\psi(\mathbf{z}) = \mathbf{\Psi} \cdot \log \mathbf{z}, \quad (13)$$

$$\mathbf{z} = \text{ilr}_\psi^{-1}(\mathbf{y}) = \mathcal{C} [\exp(\mathbf{\Psi}^- \cdot \mathbf{y})]. \quad (14)$$

Readers interested in details about these operations and transformations are referred to Tolosana-Delgado et al. (2005) where some simple cases are presented.

Seen as mappings between spaces, the log-ratio transformations all have a geometric interpretation. The component-wise logarithm $\log(\cdot)$ is a transformation from \mathbb{R}_+^D onto \mathbb{R}^D . The clr(\cdot) is a transformation from $\mathcal{S}^D \subset \mathbb{R}_+^D$ onto the hyperplane $\mathbb{H} \subset \mathbb{R}^D$ orthogonal to the vector $\mathbf{1}_D$. The matrix \mathbf{H} is thus the orthogonal projection onto this subspace. Finally, the ilr(\cdot) is a transformation from $\mathcal{S}^D \subset \mathbb{R}_+^D$ onto \mathbb{R}^{D-1} . The properties of Eq. (12) indicate that the matrix $\mathbf{\Psi}$ is an isometry on \mathbb{H} (van den Boogaart and Tolosana-Delgado, 2003).

3.3 Basic statistics

In what follows, let $\mathbf{Z} = [Z_1, Z_2, \dots, Z_D]$ denote a random composition.

Classical first and second order moments for a compositional data set can be defined through the clr or any other log-ratio transformation. If superscripts indicate the transformation used, mean and variance are given by

$$\begin{aligned}\boldsymbol{\mu}^c &= \mathbf{E}[\mathbf{H} \cdot \log(\mathbf{Z})], & \boldsymbol{\Sigma}^c &= \text{Var}[\mathbf{H} \cdot \log(\mathbf{Z})], \\ \boldsymbol{\mu}^a &= \mathbf{E}[\mathbf{J} \cdot \log(\mathbf{Z})], & \boldsymbol{\Sigma}^a &= \text{Var}[\mathbf{J} \cdot \log(\mathbf{Z})], \\ \boldsymbol{\mu}^\psi &= \mathbf{E}[\boldsymbol{\Psi} \cdot \log(\mathbf{Z})] & \boldsymbol{\Sigma}^\psi &= \text{Var}[\boldsymbol{\Psi} \cdot \log(\mathbf{Z})].\end{aligned}$$

Here the superscript c stands for clr, a for alr and ψ for an ilr transformation. Since the matrix \mathbf{H} is a projection (i.e. $\mathbf{H}^2 = \mathbf{H}$), the clr-variance matrix $\boldsymbol{\Sigma}^c$ is singular.

As an alternative to the use of a log-ratio transformation, Aitchison (1986) proposed the calculation of the *compositional centre*

$$\mathbf{m} = \mathcal{C}(\exp(\mathbf{E}[\log(\mathbf{Z})])) \quad (15)$$

and the variation matrix $\mathbf{T} = [t_{ij}]$ with elements

$$t_{ij} = \text{Var} \left[\log \frac{Z_i}{Z_j} \right] = \text{Var} [\log(Z_i) - \log(Z_j)] = \text{Var} [\text{clr}_i(\mathbf{Z}) - \text{clr}_j(\mathbf{Z})]. \quad (16)$$

One of the typical concerns of analysts when starting with the log-ratio approach to compositional data is to ensure that statistical results do not depend on the transformation used. It is well-known that the four means given before are compatible, that is (proposition 3 in the appendix)

$$\text{clr}^{-1}(\boldsymbol{\mu}^c) = \text{alr}^{-1}(\boldsymbol{\mu}^a) = \text{ilr}_\psi^{-1}(\boldsymbol{\mu}^\psi) = \mathbf{m} \in \mathcal{S}^D. \quad (17)$$

Similarly, all spread measures convey exactly the same information, and are related through the fundamental equations

$$\boldsymbol{\Sigma}^\psi = \boldsymbol{\Psi} \cdot \boldsymbol{\Sigma}^c \cdot \boldsymbol{\Psi}^t \quad \boldsymbol{\Sigma}^c = \boldsymbol{\Psi}^- \cdot \boldsymbol{\Sigma}^\psi \cdot \boldsymbol{\Psi}^{-t} \quad (18)$$

$$\boldsymbol{\Sigma}^\psi = -\frac{1}{2} \boldsymbol{\Psi} \cdot \mathbf{T} \cdot \boldsymbol{\Psi}^t \quad \boldsymbol{\Sigma}^c = -\frac{1}{2} \mathbf{H} \cdot \mathbf{T} \cdot \mathbf{H}, \quad (19)$$

where $\boldsymbol{\Psi}^{-t}$ denotes the transposed Moore-Penrose generalized inverse of $\boldsymbol{\Psi}$. Note that these expressions are valid for any basis, even a non-orthogonal one (for example an alr when taking $\boldsymbol{\Psi} = \mathbf{F}$), but at the price of having to deal with a more complex notation. Note also that this identity holds globally, for the entire composition considered, and not for individual parts. These statements are proved in, proposition 4 of the appendix.

3.4 Illustration: basic statistics

From a compositional point of view, the statistics of Table 1 are unsatisfactory, because they are not subcompositionally coherent. The set of alternative basic compositional statistics and diagrams for the K-pit data set are summarized in Table 2 and Figure 1, showing a very low variation entry for Fe-Rest, mirrored by a low-variance density plot of $\log(\text{Fe}/\text{Rest})$. This is an indication of a high proportionality between both parts, which suggests that the Rest is probably dominated by the Oxygen and OH^- -groups forming the various Fe-oxides present. In fact, the mass relation 63.2246:34.0875 on the geometric mean is almost equivalent to a 1:2 molar relation Fe:O, exactly of $(63.2246 \times 15.9994) : (34.0875 \times 55.845) = 0.53$. The kernel density estimate of $\log(\text{Fe}/\text{Rest})$ shows that most of the samples have a Fe:O molar relation between 2:5 and 2:3. Figure 1 shows very strong closure effects in the raw data (e.g., upper triangle scatterplot of Fe vs. SiO_2), but also strong boundaries on the relation between several alr-transformed variables (alr. Al_2O_3 -alr. SiO_2 , for instance), where the alr's were taken with respect to the Rest. The data are obviously not normally distributed.

Table 2: Compositional statistics of the K-Pit data set: centre (closed geometric mean) and variation matrix

	Al_2O_3	Fe	Mn	P	SiO_2	Rest
centre (%)	0.4844	63.2246	0.0344	0.1068	2.0623	34.0875
Al_2O_3	0.0000	1.0032	1.1167	1.0516	1.5451	0.8975
Fe	1.0032	0.0000	0.8876	0.3459	1.1731	0.0123
Mn	1.1167	0.8876	0.0000	0.7287	1.8846	0.8002
P	1.0516	0.3459	0.7287	0.0000	1.5046	0.3043
SiO_2	1.5451	1.1731	1.8846	1.5046	0.0000	1.1529
Rest	0.8975	0.0123	0.8002	0.3043	1.1529	0.0000

3.5 The normal distribution on the simplex

As in the case of a real variable or a real vector, it is possible to define a normally distributed random composition characterized by a given compositional center \mathbf{m} and a given variation matrix \mathbf{T} . A random composition is said to follow a normal distribution on the simplex, denoted as \mathcal{N}_{SD} , if its log-ratio transformed scores follow a joint multivariate normal distribution, i.e. with




Figure 1: Scatterplots of the K-Pit data set: raw data (upper triangle) and alr-transformed data with respect to the Rest (lower triangle). Kernel density estimates of the alr-transformed data are included in the bottom line. Note the strong effects of the closure and the positivity constraints.

probability density

$$\begin{aligned} f_{\mathbf{z}}(\mathbf{z}) &= (2\pi|\boldsymbol{\Sigma}^\psi|)^{(1-D)/2} \exp\left[-\frac{1}{2}d_{M,\mathbf{T}}^2(\mathbf{z}, \mathbf{m})\right], \\ d_{M,\mathbf{T}}^2(\mathbf{z}, \mathbf{m}) &= \text{ilr}_\psi(\mathbf{z} \ominus \mathbf{m})^t \cdot [\boldsymbol{\Sigma}^\psi]^{-1} \cdot \text{ilr}_\psi(\mathbf{z} \ominus \mathbf{m}) \end{aligned} \quad (20)$$

where the log-ratio Mahalanobis distance $d_{M,\mathbf{T}}^2(\mathbf{z}, \mathbf{m})$ denotes the log-ratio Mahalanobis distance, and $\boldsymbol{\Sigma}^\psi$ is obtained from \mathbf{T} by Eq. (19). The normal distribution on the simplex is a fundamental tool in geostatistics for compositional data, to define a Gaussian regionalized composition. Although this definition makes use of a specific ilr transformation to compute the probability density, the resulting probabilities themselves do not depend on the choice of log-ratio transformation, as long as the transformation is one-to-one. Even an alr or a clr could be used, the latter with a slightly more complicated notation. Propositions 5 and 6 in the appendix give formal proofs of these statements.

4 Conventional multivariate geostatistical practice for compositions

It is assumed here that the reader is familiar with current cokriging (simple, ordinary, universal) and cosimulation techniques, their differences and similarities (see e.g. Myers, 1982; Chilés and Delfiner, 1999; Wackernagel, 2003; Cressie, 1991) as well as with the matrix notation introduced by Myers (1982).

The focus of classical geostatistics is to estimate the value of a vector of P space-dependent (real-valued) variables $\mathbf{Z}(x) = [Z_1(x), Z_2(x), \dots, Z_P(x)]$, as a *linear function* of neighbouring observations $\{\mathbf{z}_1, \mathbf{z}_2, \dots, \mathbf{z}_N\}$ with $\mathbf{z}_i = \mathbf{Z}(x_i)$ measured at locations $\{x_1, x_2, \dots, x_N\}$ within a domain \mathcal{D} of the geographical space,

$$\phi_i(z_i^*(x_0)) = y_i^*(x) = \sum_{n=1}^N \sum_{j=1}^P w_{ij}^n \phi_j(z_j(x_n)) = \sum_{n=1}^N \sum_{j=1}^P w_{ij}^n y_j(x_n), \quad (21)$$

where and $\phi_i(\cdot), i = 1, 2, \dots, P$ can be non-linear functions: logarithms for log-normal kriging, or more typically a Gaussian anamorphosis or normal score transform (Chilés and Delfiner, 1999; Rossi and Deutsch, 2014) forcing the $y_j(x_n)$ scores to exactly show an empirical normal marginal distribution. Gaussian anamorphosis is particularly critical for simulation, as most algorithms assume the unknown scores given the available data to have a joint conditional normal distribution.

Table 3: Problems and caveats of geostatistical approaches to regionalized compositional data sets (see text for details).

method	transformation	ensures		singular kriging matrix
		$y_i^* > 0$	$\sum_i y_i^* = \kappa$	
kriging	none	no	no	no
kriging	anamorphosis	yes	no	no
	compositional kriging	yes	yes	no
cokriging	none	no	yes	yes
cokriging	anamorphosis	yes	no	no
cokriging	log-ratio	yes	yes	no

When $\mathbf{Z}(x)$ is a regionalized composition, we would like the (co)kriging systems to yield valid compositions everywhere in the domain, that is, the (co)kriging estimates satisfy the conditions stated in Eq. (1). However (as summarised in Table 3), kriging and cokriging are not convex operators (Chilés and Delfiner, 1999), thus nothing forces their results to remain bounded by the data or any constraints on them. This implies that negative predictions are possible, as are total sums larger than 100%; see e.g. Pawlowsky-Glahn et al. (1995) or Pawlowsky-Glahn and Olea (2004). Simulation algorithms do tend to generate even larger violations of these constraints, as they often add some variability around the (co)kriging predictions. Gaussian anamorphosis might ensure positivity if cleverly chosen (i.e., similar to a logarithmic transformation), but it cannot ensure the constant sum constraint. This is only ensured by cokriging (or cosimulation) of raw closed data, if one manages to deal with the singularity of the cokriging system matrix.

Methods based on normal score transforms present a secondary problem: compositional data are multivariate by nature, and some form of multivariate anamorphosis is needed. A naive marginal anamorphosis of each individual variable (as implemented in most geostatistical software) does not satisfy this condition. In this context, the stepwise conditional transform of Leuangthong and Deutsch (2003), the projection pursuit multivariate transform (PPMT Barnett et. al., 2014) or the multivariate flow normal transform of van den Boogaart et al. (2017) should be applied, as they are multivariate by construction.

Compositional kriging was introduced by Walwoort and de Gruijter (2001) to describe a cokriging of all components where positivity and constant sum constraints are included in the cokriging as additional constraints. The singularity of the covariance of raw data is dealt with by ignoring cross-covariances, thus

neglecting any natural link between variables conveyed by the cross-variograms. The only link preserved by this technique is the naive (and physically irrelevant) constant sum. Moreover, up to the authors' knowledge, this method does not deliver a meaningful cokriging covariance matrix, which makes cosimulation based on compositional kriging impossible.

Finally, a general comment on all approaches based on raw data, on simple logarithmic transformations or on individual Gaussian anamorphosis, even those taking into account the necessary constraints: These methods do not explicitly account for the relative scale of compositional data, and may thus not measure differences between observations in a sensible way. Moreover, although in some cases they satisfy numerical constraints, they do not ensure that the problem of spurious spatial covariance is avoided, which affects regionalized compositions analogously to the way in which spurious correlation affects compositions in general (Pawlowsky, 1984). To what extent these problems also affect a truly multivariate normal score transformed data set (e.g. using PPMT Barnett et. al., 2014) is still unclear.

5 Compositional structural analysis

Pawlowsky-Glahn and Olea (2004) propose several structural functions, which are essentially classical variograms and covariance functions defined either on the clr-transformed data, or on alr-transformed data. In this section we provide a summary of these definitions, and an adaptation to the case of ilr transformed data.

5.1 Structural functions and their relations

Denote by $\mathbf{Y}(x)$ the scores obtained by transforming a regionalized composition $\mathbf{Z}(x)$ through a suitable log-ratio transformation. This can be either an alr, an ilr or a clr. For any of the three transformations, a matrix-valued variogram or covariance function can be defined, as shown in proposition 9 in the appendix. The six specifications have the following properties (proposition 7):

- For a given log-ratio transformation, its variogram and covariance function satisfy

$$\mathbf{\Gamma}(h) = \mathbf{C}(0) - \frac{1}{2}(\mathbf{C}(h) + \mathbf{C}(-h));$$

thus, for a symmetric covariance function, $\mathbf{C}(h) = \mathbf{C}(-h)$, the log-ratio variogram and the log-ratio covariance function are equivalent, and they

also fulfill $\mathbf{C}(h) = \mathbf{C}(0) - \mathbf{\Gamma}(h)$. These equivalences are honored whichever log-ratio transformation is used.

- The covariance functions of ilr and clr transformed data are related by:

$$\mathbf{C}^\psi(h) = \mathbf{\Psi} \cdot \mathbf{C}^c(h) \cdot \mathbf{\Psi}^t, \quad \mathbf{C}^c(h) = \mathbf{\Psi}^t \cdot \mathbf{C}^\psi(h) \cdot \mathbf{\Psi},$$

where the superscripts c and ψ relate to the clr and ilr transformed scores, respectively.

- Analogously, the corresponding variograms are related by

$$\mathbf{\Gamma}^\psi(h) = \mathbf{\Psi} \cdot \mathbf{\Gamma}^c(h) \cdot \mathbf{\Psi}^t, \quad \mathbf{\Gamma}^c(h) = \mathbf{\Psi}^t \cdot \mathbf{\Gamma}^\psi(h) \cdot \mathbf{\Psi}. \quad (22)$$

- Equivalent expressions link clr score variograms and covariance functions with those expressed in any arbitrary basis,

$$\begin{aligned} \mathbf{C}^\psi(h) &= \mathbf{\Psi} \cdot \mathbf{C}^c(h) \cdot \mathbf{\Psi}^t, & \mathbf{C}^c(h) &= \mathbf{\Psi}^- \cdot \mathbf{C}^\psi(h) \cdot \mathbf{\Psi}^{-t}, \\ \mathbf{\Gamma}^\psi(h) &= \mathbf{\Psi} \cdot \mathbf{\Gamma}^c(h) \cdot \mathbf{\Psi}^t, & \mathbf{\Gamma}^c(h) &= \mathbf{\Psi}^- \cdot \mathbf{\Gamma}^\psi(h) \cdot \mathbf{\Psi}^{-t}. \end{aligned} \quad (23)$$

In particular, these expressions are valid when working with alr scores, by taking $\mathbf{\Psi} = \mathbf{J}$. This also applies to the remainder of this section.

The preceding relationships are satisfied both by the theoretical and the empirical versions of these functions. Thus, when fitting models to empirical variogram systems defined in two different log-ratio specifications, it is necessary to force the fitted model to satisfy them too. We say in this case that the fitted models are *mutually compatible*. In practice, ensuring compatibility during the modeling process may not be easy, as each variogram or covariance function derived from a particular log-ratio transform may focus on some specific sub-compositional features, while other might appear to mask the variability of that particular feature. It is thus possible that classical automatic fitting processes do not produce compatible models.

To avoid working with a specific transformation and having to check for mutual compatibility, a variogram function can be defined based on the variation matrix. The resulting matrix-valued function is called (pairwise) log-ratio variogram, or variation-variogram, and is denoted as the matrix $\mathbf{T}(h) = [t_{ij}(h)]_{i,j=1,\dots,D}$ with elements

$$t_{ij}(h) = \text{Var} \left[\log \frac{Z_i(x+h)}{Z_j(x+h)} - \log \frac{Z_i(x)}{Z_j(x)} \right]. \quad (24)$$

Variation-variograms contain the same information as classical clr or log-ratio coordinate variograms, so that the latter can be calculated from the former, with

$$\mathbf{\Gamma}^\psi(h) = -\frac{1}{2}\mathbf{\Psi} \cdot \mathbf{T}(h) \cdot \mathbf{\Psi}^t, \quad \mathbf{\Gamma}^c(h) = -\frac{1}{2}\mathbf{H} \cdot \mathbf{T}(h) \cdot \mathbf{H}, \quad (25)$$

and conversely from the clr-variograms

$$t_{ij}(h) = \gamma_{ii}^c(h) + \gamma_{jj}^c(h) - 2\gamma_{ij}^c(h). \quad (26)$$

There is no straightforward formula for conversion from ilr/alr-variograms to variation-variograms. Therefore, it is convenient to use Eq. (22), resp. Eq. (23), first and then Eq. (26).

5.2 Variogram estimation and modelling

Empirical variograms satisfy Eqs. (22)-(26) for all lags, if they are computed with the same set of data. Fitted models should also satisfy them. In particular, the linear model of coregionalization (LMC, Wackernagel, 2003), being symmetric, can be expressed in any of these specifications,

$$\begin{aligned} \mathbf{C}^\psi(h|\theta) &= \sum_{k=0}^K \mathbf{C}_k^\psi \cdot \rho_k(h|\theta_k), & \mathbf{C}^c(h|\theta) &= \sum_{k=0}^K \mathbf{C}_k^c \cdot \rho_k(h|\theta_k), \\ \mathbf{\Gamma}^\psi(h|\theta) &= \sum_{k=0}^K \mathbf{C}_k^\psi \cdot (1 - \rho_k(h|\theta_k)), & \mathbf{\Gamma}^c(h|\theta) &= \sum_{k=0}^K \mathbf{C}_k^c \cdot (1 - \rho_k(h|\theta_k)), \\ \mathbf{T}(h|\theta) &= \sum_{k=0}^K \mathbf{B}_k \cdot (1 - \rho_k(h|\theta_k)), \end{aligned}$$

with the vector of model parameters θ containing all individual ranges θ_k and sill matrices, specified either as \mathbf{B}_k or \mathbf{C}_k , as they are equivalent because of

$$\mathbf{C}_k^\psi = -\frac{1}{2}\mathbf{\Psi} \cdot \mathbf{B}_k \cdot \mathbf{\Psi}^t, \quad \mathbf{C}_k^c = -\frac{1}{2}\mathbf{H} \cdot \mathbf{B}_k \cdot \mathbf{H}. \quad (27)$$

A model must be positive definite if specified in terms of \mathbf{C}^ψ , and positive semi-definite in terms of \mathbf{C}^c because the clr-transformed covariances always have at least one zero eigenvalue. Equivalently, a model specification in $\mathbf{\Gamma}^\psi$ must be conditionally negative definite, and in terms of $\mathbf{\Gamma}^c$ conditionally negative semi-definite. Thus, a model specified in \mathbf{T} must be conditionally positive semi-definite. For the LMC, these conditions are satisfied if each correlogram $\rho_k(h|\theta_k)$ is a positive definite function, and each matrix \mathbf{C}_k is a valid covariance matrix (Wackernagel, 2003) or \mathbf{B}_k is a variation matrix (see the appendix), namely a conditionally negative definite matrix with zero diagonal elements.

Reduced-rank submodels of the LMC can also be used. Minimum/Maximum Autocorrelation Factors (MAF) have been applied to compositions after an alr (Morales Boezio, 2010; Morales Boezio et al., 2012; Ward and Mueller, 2012) or a clr (Mueller and Grunsky, 2016) transformation. Also, the use of rank-deficient sill matrices \mathbf{C}_k has been put forward (Gelfand and Banerjee, 2010; Tolosana-Delgado and Boogaart, 2013).

Individual variation-variograms can be estimated with any available procedure for estimation of direct variograms. Nevertheless fitting must be done jointly, to ensure that the validity conditions are satisfied. Fitting models for the variation-variograms has certain advantages. For instance it allows to use a goodness of fit that maximizes a logarithmic goodness-of-fit criterion (Tolosana-Delgado et al., 2011),

$$\text{gof}(\theta) = \sum_{i=1}^D \sum_{j=1}^D \sum_n N_{ij}(h_n) (\log \hat{t}_{ij}(h_n) - \log t_{ij}(h_n|\theta))^2, \quad (28)$$

where $\hat{t}_{ij}(h_n)$ is the empirical (i, j) -variation-variogram calculated for a lag distance class h_n , $N_{ij}(h_n)$ is the number of pairs used for that calculation, and $t_{ij}(h_n|\theta)$ is the model evaluated at lag h_n . The choice of Eq. (28) as a fit criterion focuses the fitting procedure on the smaller values of t_{ij} , i.e. around the origin and at short ranges. Two arguments support this choice. First, interpolation results are particularly sensitive to this part of the variogram, and they do not really depend on the sill: spending effort on fitting the sill will not translate in more reliable interpolations (Chilés and Delfiner, 1999). Second, as sills are variances, their natural spread ranges in orders of magnitudes, i.e. they should be compared in a relative scale, thus in logarithms: this view is consistent with the multiplicative confidence intervals given for the sill of a variogram (Cressie, 1991).

Moreover, fitting variation-variograms enables working with data sets with many missing values, partially observed subcompositions and similar irregularities. As each component $t_{ij}(h_n)$ requires only that variables i and j are available, one uses pairwise elimination to calculate $\hat{t}_{ij}(h_n)$, thus using a maximum number of observations for each lag and each pair of variables. In contrast, estimating variograms with ilr- or clr-transformed data requires eliminating much more data: for instance, the clr vector is not available if one single component is missing (i.e., one would need complete row-wise elimination to deal with missing values); and though an ilr can be chosen to maximize the number of computable log-ratio scores, this only works if the missing components are the same (Tolosana-Delgado et al., 2008).

For these reasons, a structural analysis of regionalized compositional data is better done in terms of variation-variograms. One can use standard estimation algorithms to obtain each individual pairwise log-ratio variogram. Then, a procedure is required for fitting an LMC using Eq. (28), subject to the condition that each matrix \mathbf{B}_k is a valid variation matrix (zero diagonal, conditionally negative semidefinite). Finally, the fitted model can be compared to the empirical structural functions, either in terms of the variation-variograms or recasting variogram estimates and model to clr-/ilr-variograms or covariances, using Eq. (27).

5.3 Illustration: structural analysis

The proposed compositional structural tool, the variation-variogram, is currently only available in the R package “compositions” (van den Boogaart, Tolosana-Delgado and Bren, 2009). Instead, and in order to stay within the framework of user-friendly geostatistical software, the data were alr-transformed with respect to the Rest, and the resulting scores imported into ISATIS (Geovariances, 2017). A rotation of the geographic coordinates was applied, and direct- and cross-variograms were computed at lag spacings of 2m in the vertical direction and 25m in the horizontal plane. A linear model of coregionalization (LMC) was fitted semi-automatically. The resulting parameters are summarized in Table 4 and the model fit is shown in 2.

Figures 3 and 4 show both the empirical variograms and the fitted LMC represented in variation-variograms and clr-variograms, respectively obtained with Eqs. (26) and (23). The same equations could be used to derive variograms in any isometric log-ratio representation, which were not included for space reasons. In all variogram figures, lower triangle plots (blue color) show vertical variograms, while upper triangle diagrams show (red and black/grey colors) variograms on the X-Y plane. All refer to directions after global rotation. It is worth noting that the LMC satisfactorily fits the data in all these diagrams, even though these alternative log-ratio representations were not used during the fitting process.

6 Pointwise estimation

6.1 Cokriging

Once a structural model is available, this can be used in interpolation and simulation procedures, most often ordinary cokriging. This is better done just: (a)




Figure 2: Empirical alr-variograms (dots) and fitted LMC (curves): diagonal plots (dark colors) are direct variograms, the lower triangle (blue) shows variograms in the vertical direction, the upper triangle (red, black) shows variograms in horizontal directions (after global rotation)

Table 4: Linear model of coregionalization of the alr-transformed composition;
Sph=spherical model of the indicated (X,Y,Z) ranges, after global rotation

	alr.Al ₂ O ₃	alr.Fe	alr.Mn	alr.P	alr.SiO ₂
nugget					
alr.Al ₂ O ₃	0.1324	-0.0098	0.1023	0.0415	0.0338
alr.Fe		0.0011	-0.0059	0.0018	-0.0029
alr.Mn			0.1267	0.0722	0.0177
alr.P				0.0907	-0.0072
alr.SiO ₂					0.3362
Sph(39.05, 66.94, 11.16)					
alr.Al ₂ O ₃	0.225	-0.0026	-0.02	-0.0042	0.0253
alr.Fe		0.0034	-0.0033	-0.0033	-0.0033
alr.Mn			0.1575	0.0138	-0.0650
alr.P				0.0554	0.0145
alr.SiO ₂					0.1330
Sph(3.3·10 ² 2.0·10 ² 1.0·10 ⁸)					
alr.Al ₂ O ₃	0.3901	-0.0147	0.0364	-0.0602	0.1710
alr.Fe		0.0043	-0.0059	0.0056	-0.0134
alr.Mn			0.2806	0.0515	0.0685
alr.P				0.1159	-0.0389
alr.SiO ₂					0.3812
Sph(34.8, 34.8, 34.8)					
alr.Al ₂ O ₃	0.199	-0.0118	0.0864	0.0586	0.1185
alr.Fe		0.003	-0.0109	-0.0147	0.0073
alr.Mn			0.1998	0.042	-0.0045
alr.P				0.0753	-0.0409
alr.SiO ₂					0.3486




Figure 3: Empirical variation-variograms (dots) and fitted LMC (curves): the lower triangle (blue) shows variograms for the vertical direction, the upper triangle (red, black) variograms in horizontal directions (after global rotation).




Figure 4: Empirical clr-variograms (dots) and fitted LMC (curves): diagonal plots (dark colors) are direct variograms, the lower triangle (blue) shows variograms in the vertical direction, the upper triangle (red, black) show variograms in horizontal directions (after global rotation)

choosing a particular ilr coordinate system, (b) expressing data and variation-variogram model in this system by Eqs. (13) and (27) respectively, (c) applying cokriging/cosimulation to the obtained log-ratio scores, and (d) back-transforming the interpolated/cosimulated scores to interpolated/cosimulated compositions through Eq. (14). This procedure ensures that the final outcomes are exactly the same, whichever log-ratio transform was used (proposition 9). It is nevertheless recommended to avoid the clr transformation, as clr scores sum to zero, and this collinearity makes the cokriging matrix singular. This could be resolved by Moore Penrose generalized inversion, if this is available, but producing the same solution at a larger computational cost is seldom a good choice.

6.2 Cross-validation and error assessment

When faced with the application of log-ratio methods, practitioners often wonder whether this approach is “better” or “worse” than “classical” methods, and try to answer that by a cross-validation exercise, where a root mean square error (RMSE), a standardized residual sum of squares (STRESS) or a similar measure of difference between observations and independent predictions is drawn (Lark and Bishop, 2007; Morales Boezio, 2010; Sun et al., 2014; Rossi and Deutsch, 2014). It should be noted that this procedure is fair only when comparing methods that minimize the same error measure, it is not appropriate to compare estimates obtained with Eq. (21) by using two different $\phi(\cdot)$ functions, e.g. the identity (cokriging raw data) vs. a log-ratio transformation (cokriging log-ratio data).

If the goodness-of-fit measure is related to a distance $d(\mathbf{y}(x_i), \hat{\mathbf{y}}_i)$, then estimators minimizing its expected value will be favored. Pawłowsky-Glahn and Egozcue (2001) called such estimators *metric expectations*, and Pawłowsky-Glahn and Egozcue (2002) showed that these coincide with linear estimators in terms of any isometric transformation of the data (isometric with respect to the distance chosen). As a consequence, if the chosen measure of goodness of fit is an absolute difference between estimates and true values, one immediately favors cokriging of raw components. In contrast, by choosing a relative difference (like the Aitchison distance), one favors log-ratio methods. There is thus no fair comparison between them.

A second consideration relates to the multivariate nature of compositional data. A compositional goodness-of-fit cross-validation criterion should be multivariate rather than univariate. For instance, one could compare some distance

between the observed compositions $\mathbf{y}(x_i)$ and their cross-validation predictions $\hat{\mathbf{y}}_i$, preferentially a Mahalanobis distance (Eq. 20) taking into account the cokriging error covariance (Lark and Bishop, 2007).

Connecting these two ideas, it seems reasonable to evaluate compositional methods by a Mahalanobis distance between predictions and values of the log-ratio transformed data (20). Note that this Mahalanobis distance will deliver the same results, whichever log-ratio transform is used, as long as it is one-to-one (Filzmoser and Hron, 2008, or Proposition 5 in the appendix). This property is not satisfied by the Euclidean distance of the scores, which requires working with isometric transformations (clr or ilr) in order to give rise to the Aitchison distance (Eq. 6).

Classical cross-validation (leave-one-out, n-fold, etc.) typically involves a visual assessment of the adequacy of a series of diagrams, judged by experience. These diagrams are histograms of residuals or standardized residuals, scatter plots of residuals against predictions, and scatterplots of true values against predictions (Bivand et. al., 2013). For the purposes of validation with compositional data, all these scatter plots can be obtained for the log-ratio scores; and for the back-transformed components as well, if the input data and the cross-validated values are computed in the same (sub)composition. Histograms and qq-plots, on the other hand, are better obtained only for the log-ratio scores, because the original components will have strongly non-normal distributions. Histograms of Aitchison-Mahalanobis norms of residuals may also be useful.

6.3 Illustration: cokriging and cross-validation

Using the LMC of section 5.3, a 5-fold cross-validation was conducted to assess the adequacy of the LMC. The data set was randomly split into 5 subsets, and for each subset co-kriging estimates were computed based on the data from the remaining subsets. Cokriging covariance matrices were also produced. Computations were obtained with R. Figure 5 shows the estimates against true values and against residuals in log-ratio scale, while figure 6 shows them in raw scale, i.e. in the original units. In all cases, a red line shows a reference line, be it the identity line for plots of observations vs predictions, or the horizontal line $y = 0$ for diagrams of residuals vs predictions. All diagrams show a reasonable absence of trends or non-linearity structures, although some residuals in raw scale show a clear heteroskedasticity. Adjustment of the data to the reference lines is also very good.

As a multivariate measure of goodness of fit, one can study the Aitchison-




Figure 6: Scatter density of residuals against predictions (upper row) and of true values against predictions (lower row) for a 5-fold cross-validation. Each diagram represents outcomes for one original component, in %. Regression lines (red) and correlation coefficients are reported. Reference lines are also shown in black: the $y = 0$ line for the upper row plots, the $y = X$ line for the lower row plots.




Figure 7: Comparison of Aitchison-Mahalanobis cokriging residual norms with a χ^2 -distribution of 5 degrees of freedom (in log scale): histogram (left) and qq-plot (right)

Mahalanobis cokriging residual (squared) norms: these are (squared) Aitchison-Mahalanobis distances (Eq. 20) between true values and cokriging estimates, with respect to the cokriging covariance matrix. Given that the dimension of the space is 5, these squared norms should follow a χ^2 distribution with 5 degrees of freedom. Figure 7 shows this comparison, in the form of a histogram and a qq-plot for a robust subset of the predictions. This selection was necessary because relative residuals are non-robust quantities: to filter the data exhibiting this problem, it is common in conventional geostatistical practice to filter standardized residuals larger than 2.5 in absolute value. The outcome shows a reasonable fit, although with a slightly heavier tail.

7 Simulation and blockwise estimation

7.1 Conditional joint distribution

As happens with conventional (real-valued) variables, cokriging of regionalized compositional data sets delivers something more than just an “interpolation”. Ordinary cokriging provides the expected value of the ilr-transformed composition at the predicted location given the surrounding observed compositions. And it can, by virtue of Eq. (17), be back-transformed to an unbiased estimate of the corresponding conditional expected value of the composition. This unbiasedness is with respect to the Euclidean structure introduced in Section 3.2 (Pawlowsky-Glahn and Egozcue, 2002). Moreover, under the key assumption of

(transformed) Gaussianity discussed in the Introduction, simple cokriging delivers the full conditional distribution of the ilr-transformed composition at the predicted location given the surrounding observed compositions: a multivariate normal distribution with mean equal to the simple cokriging predictor and covariance matrix equal to the cokriging error covariance. But as stated in Section 3.5, these two moments of any ilr representation completely specify the normal distribution on the simplex that describes the conditional uncertainty on the interpolated composition. Conditional distributions are primarily used for two purposes: cosimulation and estimation of non-linear quantities.

7.2 Cosimulation

Any cosimulation algorithm applied to the log-ratio transformed composition will provide valid simulated log-ratios, that can be back transformed to simulated compositions through Eqs. (7-13). Mueller et al. (2014) show that this procedure adequately reproduces the mean value and the sample distribution of the variables (on both scales: the original proportions and the log-ratio transformed values), as long as the log-ratio transformed data do not depart notably from normality. Reproduction of the variation-variogram is, by construction, as good (or as bad) as the simulation method actually used. These caveats are well-known for simulation procedures of raw data, which often rely on Gaussian anamorphosis to enforce these global statistics reproduction. Recent concerns (Rossi and Deutsch, 2014) on this side for the log-ratio procedure are thus quite partial.

As a matter of good practice, one should avoid any procedure for which final simulated compositions would depend on the log-ratio transformation used. For this reason, separate simulation of each log-ratio variable (with or without a normal score transform) should be avoided. However, a multivariate Gaussian anamorphosis (Barnett et. al., 2014) will be necessary when departures from additive logistic normality are notable. The recently introduced flow anamorphosis (van den Boogaart et al., 2017) provides a Gaussian anamorphosis that ensures independence of the results from the log-ratio transform applied.

7.3 Estimation of non-linear quantities

With regard to the estimation of non-linear quantities (non-linear in terms of the log-ratio transformed composition), either Monte Carlo simulation or Gauss-Hermite quadratures can be used. If \hat{y}_0 and $\hat{\mathbf{S}}_y$ denote respectively the simple cokriging estimate and covariance matrix of an ilr-transformed composition

$\mathbf{Y}(x_0)$, then

$$[\mathbf{Y}(x_0)|\mathbf{y}_1, \dots, \mathbf{y}_N] \sim \mathcal{N}^{D-1}(\hat{\mathbf{y}}_0, \hat{\mathbf{S}}_y). \quad (29)$$

If interest lies in a non-linear function $g(\mathbf{Y}(x_0))$, a Monte Carlo sample of this quantity can be obtained from a set of K simulations $\{\tilde{\mathbf{y}}^{(1)}, \dots, \tilde{\mathbf{y}}^{(K)}\}$. By transforming each one of them, the set $\{g(\tilde{\mathbf{y}}^{(1)}), \dots, g(\tilde{\mathbf{y}}^{(K)})\}$ will describe the distribution of $g(\mathbf{Y}(x_0))$, and any relevant statistics (mean, variance, quantiles, etc.) can be derived from this transformed sample. This Monte Carlo strategy can also be used for spatially averaged quantities, such as block kriging estimates (Tolosana-Delgado et al., 2013).

An alternative is the use of Gauss-Hermite quadratures to estimate the expected value of $g(\mathbf{Y}(x_0))$. This requires a factorization of $\hat{\mathbf{S}}_y = \mathbf{R}\mathbf{R}^t$, e.g. a Cholesky decomposition or a decomposition based on the singular value decomposition of $\hat{\mathbf{S}}_y$. Applying the change of variable $\mathbf{U} = \mathbf{R}^{-1} \cdot (\mathbf{Y} - \hat{\mathbf{y}}_0)/\sqrt{2}$, the expectation sought can be written as

$$\int_{\mathbb{R}^{D-1}} g(\mathbf{Y})\phi(\mathbf{Y}|\hat{\mathbf{y}}_0, \hat{\mathbf{S}}_y)d\mathbf{Y} = \int_{\mathbb{R}^{D-1}} \pi^{-(D-1)/2} g(\mathbf{Y}(\mathbf{U})) \exp(-\mathbf{U}^t\mathbf{U})d\mathbf{U} = I.$$

An order k Gauss-Hermite approximation can be computed using weights w_1, w_2, \dots, w_k and quadrature points u_1, u_2, \dots, u_k as

$$I \approx \sum_{i_1=1}^k \sum_{i_2=1}^k \cdots \sum_{i_{D-1}=1}^k w_{i_1} w_{i_2} \cdots w_{i_{D-1}} g(\hat{\mathbf{y}}_0 + \sqrt{2} \cdot \mathbf{R} \cdot \mathbf{u}_{[i_1, i_2, \dots, i_{D-1}]}) \quad (30)$$

involving k^{D-1} $(D-1)$ -tuples of Hermite quadrature points $\mathbf{u}_{[i_1, i_2, \dots, i_{D-1}]} = [u_{i_1}, u_{i_2}, \dots, u_{i_{D-1}}]$. These expressions were used by Aitchison (1986, p. 314) to obtain estimates of the composition $\mathbf{Z}(x_0)$ unbiased in terms of the original units (ppm, %, etc) under the assumption that the geometry of real space induced in the simplex holds. This is readily available using the vector-valued function $g(\mathbf{Y}(x_0)) = \text{ilr}^{-1}(\mathbf{Y}(x_0))$ (Pawlowsky-Glahn and Olea, 2004; Lark and Bishop, 2007; Ward and Mueller, 2013). Note that the final results of this approximation to I might depend on which log-ratio was used (see the appendix). Being a Gauss quadrature approximation of $E[\mathbf{Z}(x_0)]$, the result of this numerical integration is an unbiased estimator of $\mathbf{Z}(x_0)$ in the classical, Euclidean difference based sense.

7.4 Illustration: cosimulation in point and block support

Cosimulation requires joint normality of the log-ratio scores. Figure 1 (lower triangle plots) show that the alr data do not have uni- and bi- normal marginals

(nor will any other log-ratio transform). Hence a form of joint multivariate normal anamorphosis is required. We applied the flow anamorphosis (van den Boogaart et al., 2017), with parameters $\sigma_0 = 0.1$ and $\sigma_1 = 1.1$. Resulting scores were shown to be spatially uncorrelated, something often observed in flow-anamorphosed scores. Thus, only direct variograms of the 5 normal scores were modelled, with structures reported in Table 5. This allowed applying a separate simulation procedure for each of them. In a general framework with non-zero cross-variograms, one should use cosimulation in order to ensure that results are independent of the logratio transform used.

Table 5: Variogram models of the flow-anamorphosed scores; Sph=spherical model of the indicated (X,Y,Z) ranges, after global rotation

	Ana ₁	Ana ₂	Ana ₃	Ana ₄	Ana ₅
Nugget	0.24	0.18	0.26	0.13	0.13
Sill ₁	0.22	0.25	0.15	0.26	0.3
Sph ₁	(20, 5, 24)	(20, 5, 24)	(40, 5, 20)	(14, 14, 5)	(14, 20, 10)
Sill ₂	0.19	0.19	0.3	0.31	0.35
Sph ₂	(60, 100, 14.5)	(30, 100, 14.5)	(50, 70, 28)	(60, 60, 30)	(200, 100, 30)
Sill ₃	0.36	0.38	0.3	0.3	0.22
Sph ₃	(600, 200, ∞)	(400, 200, ∞)	(200, 400, ∞)	(180, 120, ∞)	(200, 300, ∞)

The turning bands algorithm in ISATIS was used to generate the realisations. Outcomes were backtransformed via the inverse flow anamorphosis, and the inverse alr-transform to obtain point-support simulations of the composition. These were upscaled to block support, with blocks of $12 \times 12 \times 6\text{m}^3$, that is, averaging simulations at 8 points. Figure 9 shows two views of the results. The spatial averages were all computed in mass scale, that is adding tons/block of component i at the different locations.

The E-type estimates were computed on the relevant scale. For single variables expressed in percentages, the relevant scale is additive, because the goal is to generate unbiased estimates of the mass of a certain element within each block, and masses are additive. In contrast, estimates of proportions should be unbiased regardless of what subcomposition is considered, i.e. a compositional scale is relevant. Hence, out of $N_{sim} = 100$ simulations, E-types for %Fe at block V were obtained as

$$Etype[\%Fe(V)] = \frac{1}{N_{sim}} \sum_{k=1}^{N_{sim}} \sum_{x \in V} w^{(k)}(x) Fe^{(k)}(x), \quad (31)$$




Figure 8: Experimental variograms and fitted models for the flow-anamorphosed variables. The top row (red, black) shows direct variograms in horizontal directions (after global rotation), the bottom row (blue) those in the vertical direction.

while E-types for the (log)-proportion Fe:Mn were produced with

$$Etype \left[\log \frac{Fe(V)}{Mn(V)} \right] = \frac{1}{N_{sim}} \sum_{x \in V} \log \frac{\sum_{x \in V} w^{(k)}(x) Fe^{(k)}(x)}{\sum_{x \in V} w^{(k)}(x) Mn^{(k)}(x)}, \quad (32)$$

where $Fe^{(k)}(x)$ or $Mn^{(k)}(x)$ denote the values of iron or manganese at a location x for the k -th simulation, and $w^{(k)}(x)$ is the tonnage value allocated to point x at simulation k . This is the portion of mass allocated to a small block v of $6 \times 6 \times 3\text{m}^3$ centered at x with respect to the mass of the block V of $12 \times 12 \times 6\text{m}^3$. Estimating it properly requires cosimulating the rock density $\rho(x)$ as well (Tolosana-Delgado et al., 2015). In this illustration the density was assumed constant, so that $w^{(k)}(x) = 1/8$ for this configuration of simulation points within the block V . Eq. (32) assumes that the processing of the block will occur after some form of spatial mixing, hence the spatial average is taken as an arithmetic mean: this will be a common situation in mining, due to the effect of milling. However, one can imagine other situations in which a spatial average is not relevant, in which case a quantity such as

$$Etype \left[\log \frac{Fe}{Mn}(V) \right] = \frac{1}{N_{sim}} \sum_{x \in V} w^{(k)}(x) \log \frac{Fe^{(k)}(x)}{Mn^{(k)}(x)}, \quad (33)$$

would be more appropriate.




Figure 9: North view of the deposit, with indication of block %Fe (top) and of $\log(Fe/Mn)$ (bottom), both E-type estimates out of 100 simulations. Two horizontal slices, 7 vertical transversal slices and one vertical longitudinal slice of the block model are shown.

8 Conclusions

Compositional data should be treated geostatistically expressed in terms of log-ratios. The idea is to use conventional multivariate geostatistical tools in the transformed scores, and back-transform interpolations or simulations. This is analogous to the conventional gaussian anamorphosis or normal score transform, but multivariate and concept-driven, instead of univariate and data-driven.

All log-ratio transformations available yield the same final compositional results, as long as the transformation is one-to-one and the variogram models used are compatible with each other. Indeed, given that the choice of one or another log-ratio transformation is somewhat an arbitrary one, this invariance (*affine equivariance*) is highly desirable. Moreover, it makes to discuss or consider that each *alr* or *ilr* could give different results. However, numerical approximations of non-linear transformations (for instance, the unbiased estimations of the composition in percentages within the original subcomposition) obtained by Gauss-Hermite quadratures might differ because of the approximation error. Note that no univariate normal score transformations (Gaussian namorphosis) is admissible, if this invariance is desired. Appropriate multivariate Gaussian anamorphoses exist, like the flow anamorphosis, which is by construction affine equivariant.

The variographic structure should be characterized by exploring several possible log-ratio sets, to check that the fit of the model to the empirical variograms is consistent across them. These should include at least variation-variograms, formed by direct variograms of all pairwise log-ratios. in addition to classical automatic fitting procedures, these variation-variograms can be fitted with a logarithmic goodness of fit criterion which focuses on improving the fit at short ranges.

Finally, bias and cross-validation error measures should be computed with caution. First, they should not be used at all to rank methods that minimize different kriging error variances. Second, given the multivariate nature of compositional data, multivariate measures of goodness-of-fit should be preferred over an individual component by component error assessment: a distance (preferentially an Aitchison-Mahalanobis distance) between each vector of observations and its predictions should be used.

With regard to representating compositions, a powerful option is to use matrices of pairwise log-ratios. This concept was first used in the variation matrix, as a matrix of variances, and has been extended to diagrams like the variation-variogram, or the visualization of cross-validation prediction vs true

values.

Upscaling to block estimates as well as computing expectations of non-linear functions of the composition can be achieved by using simulations. This typically requires taking a multivariate Gaussian anamorphosis of the logratio transformed scores, and apply conventional variography and cosimulation procedures to the Gaussian scores obtained. Simulations can then be back-transformed to logratio scores and to compositions, and then upscaled or transformed as required. An average of the results will deliver a Monte Carlo estimation of the target quantities.

A Appendix: Formal statements and proofs

In this appendix, proofs for the results in the main text are provided. They are organised into a “non-spatial” part and a spatial part. In what follows the variables \mathbf{z} and \mathbf{Z} will denote the composition on the original scale and $\boldsymbol{\zeta}$ or \mathbf{Z} will be used for the log-ratio transformed data.

A.1 Non-spatial results

Definition 1 (Composition as closed vectors) *A vector $\mathbf{z} \in \mathbb{R}^D$ is called a composition if its k^{th} component z_k represents the relative importance of part k with respect to the remaining components.*

Typically, $z_k \geq 0$ and $z_1 + z_2 + \dots + z_D = \kappa$, with $\kappa = 1$ (for proportions), $\kappa = 100$ (for percentages) and $\kappa = 10^6$ (for ppm). However, the variables under consideration might only represent a subset of all possible variables in which case the constant sum constraint is not necessarily satisfied. Subsequent treatment of the data then depends on whether or not the resulting non-constant sum is meaningful and less than κ . In this case a fill-in variable (Eq. 2) can be added to retain that information and fulfill the constraint. On the other hand, if the non-constant sum is meaningless, the data can be reclosed (Eq. 3) without losing any information. Mathematically, this last case gives rise to the definition of compositions as equivalence classes (Barceló-Vidal, 2003), the modern, more general definition of composition.

Definition 2 (log-ratio representation) *A function $\psi(\cdot)$ is a full-rank log-ratio representation of the composition \mathbf{z} if its image satisfies*

$$\psi(\mathbf{z}) = \boldsymbol{\Psi} \cdot \log \mathbf{z} =: \boldsymbol{\zeta}$$

where $\boldsymbol{\Psi}$ is a $(D - 1) \times D$ matrix of rank $(D - 1)$ with $\boldsymbol{\Psi} \cdot \mathbf{1} = \mathbf{0}$.

Lemma 1 (inversion) *If $\psi(\cdot)$ is a full-rank log-ratio transformation, then the corresponding matrix Ψ satisfies $\Psi^- \cdot \Psi = \mathbf{H}$, where \mathbf{H} is the projection on the orthogonal complement of the vector $\mathbf{1}$ in \mathbb{R}^D .*

Proof: The singular value decomposition of Ψ is given by $\Psi = \mathbf{U} \cdot \mathbf{S} \cdot \mathbf{V}^t$, where \mathbf{U} is an orthogonal $(D-1) \times (D-1)$ matrix, \mathbf{V} is an orthogonal $D \times D$ matrix with $\mathbf{V}^t \mathbf{V} = \mathbf{I}_{(D-1)}$ and $\mathbf{S} = \begin{bmatrix} \mathbf{D}_{(D-1)} & \mathbf{0}_{(D-1)} \end{bmatrix}$ is a $(D-1) \times D$ matrix with \mathbf{D} an invertible real diagonal matrix and $\mathbf{0}$ a column vector of zeros. The Moore-Penrose inverse is therefore $\Psi^- = \mathbf{V} \cdot \mathbf{D}^+ \cdot \mathbf{U}^t$ where $\mathbf{D}^+ = [\mathbf{S}^{-1} \ \mathbf{0}]'$. Then

$$\begin{aligned} \Psi^- \cdot \Psi &= (\mathbf{V} \cdot \mathbf{S}^- \cdot \mathbf{U}^t) \cdot (\mathbf{U} \cdot \mathbf{S} \cdot \mathbf{V}^t) \\ &= \mathbf{V} \cdot \begin{bmatrix} \mathbf{I}_{(D-1)} & \mathbf{0} \\ \mathbf{0}^t & 0 \end{bmatrix} \cdot \mathbf{V}^t. \end{aligned}$$

Since $\mathbf{D}^+ \cdot \mathbf{D}$ has rank $D-1$ and \mathbf{V} has full rank, $\Psi^- \cdot \Psi$ has rank $D-1$ and its eigenvalues are 1 and 0. Since $(\Psi^- \cdot \Psi)^2 = \Psi^- \cdot \Psi$, and $(\Psi^- \cdot \Psi)^t = \Psi^- \cdot \Psi$, the matrix $\Psi^- \cdot \Psi$ is an orthogonal projection. Moreover from the definition of Ψ it follows that $\Psi^- \cdot \Psi \cdot \mathbf{1} = \Psi^- \cdot \mathbf{0} = \mathbf{0}$. Therefore, if the columns of \mathbf{V} are denoted by $\mathbf{v}_i, i = 1, \dots, D$, the eigenvector for 0 is given by $\mathbf{v}_D = \frac{1}{\sqrt{D}} \mathbf{1}$, so that $(\Psi^- \cdot \Psi) = \mathbf{I}_D - \mathbf{v}_D \mathbf{v}_D^t = \mathbf{I}_D - \frac{1}{D} \mathbf{1}_{D \times D} = \mathbf{H}$. \square

Proposition 1 (Inverse log-ratio representation) *A full-rank log-ratio representation $\psi(\cdot)$ is one-to-one, and its inverse is*

$$\mathbf{z} = \mathcal{C}[\exp(\Psi^- \cdot \zeta)].$$

Proof: From the previous lemma it follows that

$$\begin{aligned} \mathcal{C}[\exp(\Psi^- \cdot \zeta)] &= \mathcal{C}[\exp(\Psi^- \cdot \Psi \cdot \log \mathbf{z})] \\ &= \mathcal{C}[\exp((\mathbf{H} \cdot \log \mathbf{z})] \\ &= \mathcal{C}[\exp(\text{clr}(\mathbf{z}))] \equiv \mathbf{z}. \end{aligned}$$

It remains to be shown that $\psi(\cdot)$ is one-to-one when restricted to the orthogonal complement of $\mathbf{1}_D$, but this is a direct consequence of the definition of $\psi(\cdot)$. \square

Proposition 2 (Change of log-ratio representation) *Let \mathbf{z} be a composition, and $\psi_1(\cdot)$ and $\psi_2(\cdot)$ be two full-rank log-ratio transformations characterized by the matrices Ψ_1 and Ψ_2 respectively. Then, its two log-ratio representations $\zeta_1 = \psi_1(\mathbf{z})$ and $\zeta_2 = \psi_2(\mathbf{z})$ are related through the linear relationship*

$$\zeta_2 = \mathbf{A}_{12} \cdot \zeta_1 \tag{34}$$

where the matrix $\mathbf{A}_{12} = \Psi_2 \cdot \Psi_1^-$ is square and invertible.

Proof: From the preceding two propositions it follows that $\zeta_2 = \psi_2(\mathbf{z}) = \Psi_2 \cdot \log \mathbf{z}$ and $\mathbf{z} = \mathcal{C}[\exp(\Psi_1^- \cdot \zeta_1)]$. Substituting the second expression into the first,

$$\zeta_2 = \Psi_2 \cdot \log (\mathcal{C}[\exp(\Psi_1^- \cdot \zeta_1)]) = \Psi_2 \cdot [\Psi_1^- \cdot \zeta_1 - \alpha \mathbf{1}] = \Psi_2 \cdot \Psi_1^- \cdot \zeta_1,$$

where $\alpha = \log(\mathbf{1}^t \cdot \exp(\Psi_1^- \cdot \zeta_1))$. This last term satisfies $\kappa \Psi_2 \cdot \mathbf{1} = \mathbf{0}$, which delivers the final expression as sought. \square

Proposition 3 (log-ratio representation of the mean) *Let $\mathbf{Z} = [z_{kn}]$, $k = 1, 2, \dots, D$, $n = 1, 2, \dots, N$, be a compositional data set with N observations and D parts, and $\psi(\cdot)$ be a full-rank log-ratio transformation. Then $\hat{\mathbf{E}}[\psi(\mathbf{Z})] = \psi(\hat{\mathbf{m}})$ the log-ratio representation of the closed geometric mean (Eq. 15).*

Proof: The empirical closed geometric center is $\hat{\mathbf{m}} = \mathcal{C}[\exp(\log(\mathbf{Z}) \cdot \mathbf{1}_N/N)]$. The log-ratio mean is given by $\hat{\mathbf{E}}[\psi(\mathbf{Z})] = (\Psi \cdot \log \mathbf{Z}) \cdot \mathbf{1}_N/N$. Substituting this expression into the definition of the inverse log-ratio representation results in

$$\begin{aligned} \psi^{-1}(\hat{\mathbf{E}}[\psi(\mathbf{Z})]) &= \mathcal{C}[\exp(\Psi^- \cdot \hat{\mathbf{E}}[\psi(\mathbf{Z})])] = \mathcal{C}[\exp(\Psi^- \cdot \Psi \cdot \log(\mathbf{Z}) \cdot \mathbf{1}_N/N)] = \\ &= \mathcal{C}[\exp(\log(\mathbf{Z}) \cdot \mathbf{1}_N/N)] = \hat{\mathbf{m}}. \end{aligned}$$

\square

This proposition also proves Eq. (17): Because the calculation of \mathbf{m} does not involve any log-ratio representation, all log-ratio representations are equivalent.

Proposition 4 (log-ratio representations of the covariance) *Let $\mathbf{Z} = [z_{kn}]$, $k = 1, 2, \dots, D$, $n = 1, 2, \dots, N$, be a compositional data set with N observations and D parts, and $\psi(\cdot)$ be a full-rank log-ratio transformation. Then the covariance matrix of the log-ratio representation can be obtained from the empirical variation matrix $\hat{\mathbf{T}}$ as $\hat{\Sigma}^\psi = -\frac{1}{2} \Psi \cdot \hat{\mathbf{T}} \cdot \Psi^t$.*

Proof: From (Aitchison, 1986) it is known that the clr covariance $\hat{\Sigma}^c$ is related to the empirical variation matrix by $\hat{\Sigma}^c = -\frac{1}{2} \mathbf{H} \cdot \hat{\mathbf{T}} \cdot \mathbf{H}$ and $\Psi \cdot \mathbf{H} = \Psi$, which is a consequence the definition of the matrix \mathbf{H} (Eq 9),

$$\Psi \cdot \mathbf{H} = \Psi \cdot \left(\mathbf{I}_{D \times D} - \frac{1}{D} \mathbf{1}_{D \times D} \right) = \Psi \cdot \mathbf{I}_{D \times D} - \frac{1}{D} \Psi \mathbf{1}_{D \times D} = \Psi - \frac{1}{D} \mathbf{0} = \Psi,$$

because the rows of Ψ sum to zero. Therefore it remains to be shown $\hat{\Sigma}^\psi = \Psi \cdot \hat{\Sigma}^c \cdot \Psi^t$. The (maximum likelihood) estimators of these two covariance matrices are

$$\begin{aligned} \hat{\Sigma}^\psi &= \frac{1}{N} (\Psi \cdot (\log(\mathbf{Z}) - \log(\hat{\mathbf{m}}) \cdot \mathbf{1}_N^t)) \cdot ((\log(\mathbf{Z}) - \log(\hat{\mathbf{m}}) \cdot \mathbf{1}_N^t)^t \cdot \Psi^t) \\ \hat{\Sigma}^c &= \frac{1}{N} (\mathbf{H} \cdot (\log(\mathbf{Z}) - \log(\hat{\mathbf{m}}) \cdot \mathbf{1}_N^t)) \cdot ((\log(\mathbf{Z}) - \log(\hat{\mathbf{m}}) \cdot \mathbf{1}_N^t)^t \cdot \mathbf{H}). \end{aligned}$$

Since $\mathbf{H} = \mathbf{H}^t$, so that $\mathbf{H} \cdot \Psi^t = \Psi^t$, it follows that

$$\begin{aligned}\hat{\Sigma}^\psi &= \frac{1}{N} (\Psi \cdot \mathbf{H} \cdot (\log(\mathbf{Z}) - \log(\hat{\mathbf{m}}) \cdot \mathbf{1}_N^t)) \cdot ((\log(\mathbf{Z}) - \log(\hat{\mathbf{m}}) \cdot \mathbf{1}_N^t)^t \cdot \mathbf{H} \cdot \Psi^t) \\ &= \Psi \cdot \hat{\Sigma}^c \cdot \Psi^t.\end{aligned}$$

Therefore $\hat{\Sigma}^\psi = \Psi \cdot \hat{\Sigma}^c \cdot \Psi^t = -\frac{1}{2} \Psi \cdot \mathbf{H} \cdot \hat{\mathbf{T}} \cdot \mathbf{H} \cdot \Psi^t = -\frac{1}{2} \Psi \cdot \hat{\mathbf{T}} \cdot \Psi^t$ \square

It is straightforward to show that the same properties hold for unbiased estimators (with denominator $N - 1$).

The preceding two propositions show that the empirical log-ratio mean vector and covariance matrix can be obtained directly from the empirical closed geometric center and variation matrix. Equivalent relationships exist also between the theoretical counterparts of these statistics.

Corollary 1 *If $\psi(\cdot)$ is a full rank log-ratio transform, then $\Psi^- \cdot \hat{\Sigma}^\Psi \cdot \Psi^{-t} = \hat{\Sigma}^c$.*

Proof: From Proposition 4 it follows that $\Psi^- \cdot \hat{\Sigma}^\Psi \cdot \Psi^{-t} = \Psi^- \cdot \Psi \cdot \hat{\Sigma}^c \cdot \Psi^t \cdot \Psi^{-t} = \mathbf{H} \cdot \hat{\Sigma}^c \cdot \mathbf{H}^t = \hat{\Sigma}^c$. \square

Corollary 2 *If $\psi_1(\cdot)$ and $\psi_2(\cdot)$ are full rank log-ratio transforms, then $\hat{\Sigma}^{\psi_2} = \mathbf{A}_{12} \cdot \hat{\Sigma}^{\psi_1} \cdot \mathbf{A}_{12}^t$.*

Proof: From $\hat{\Sigma}^{\psi_2} = \Psi_2 \cdot \hat{\Sigma}^c \cdot \Psi_2^t$ and Corollary 1 it follows that $\hat{\Sigma}^{\psi_2} = \Psi_2 \cdot \Psi_1^- \cdot \hat{\Sigma}^{\psi_1} \cdot \Psi_1^{-t} \cdot \Psi_2^t = \mathbf{A}_{12} \cdot \hat{\Sigma}^{\psi_1} \cdot \mathbf{A}_{12}^t$. \square

Corollary 3 *If $\psi(\cdot)$ is a full rank log-ratio transform, then $\Sigma^{c-} = \Psi^t \cdot \Sigma^{\psi^{-1}} \cdot \Psi$ is a generalised inverse of Σ^c .*

Proof: Firstly, Σ^Ψ has full rank and so is invertible, thus

$$\begin{aligned}\Sigma^c \cdot \Sigma^{c-} &= \Psi^- \cdot \Sigma^\psi \cdot \Psi^{-t} \Psi^t \cdot (\Sigma^\psi)^{-1} \cdot \Psi \\ &= \Psi^- \cdot \Sigma^\psi \cdot (\Psi \cdot \Psi^-)^t \cdot (\Sigma^\psi)^{-1} \cdot \Psi \\ &= \Psi^- \cdot \Sigma^\psi \Sigma^{\psi^{-1}} \cdot \Psi \\ &= \Psi^- \Psi = \mathbf{H}.\end{aligned}$$

since $(\Psi \cdot \Psi^-) = \mathbf{I}_{(D-1)}$, so that $\Sigma^c \cdot \Sigma^{c-}$ is symmetric. Secondly, $\Sigma^c \cdot \Sigma^{c-} \cdot \Sigma^c = \mathbf{H} \cdot \Sigma^c = \Sigma^c$ and $\Sigma^{c-} \cdot \Sigma^c \cdot \Sigma^{c-} = \Sigma^{c-} \cdot \mathbf{H} = \Psi^t \cdot (\Sigma^\psi)^{-1} \cdot \Psi \cdot \mathbf{H} = \Sigma^{c-}$. Similarly, $\Sigma^{c-} \cdot \Sigma^c = \mathbf{H}$. Therefore $\Psi^t \cdot (\Sigma^\psi)^{-1} \cdot \Psi$ satisfies all conditions of a generalised inverse. \square

Proposition 5 (Invariance of the Mahalanobis distance) *Let \mathbf{Z} be a composition, with variation matrix \mathbf{T} . The Aitchison-Mahalanobis distance between any two of its realizations \mathbf{z}_1 and \mathbf{z}_2*

$$d_{M,\mathbf{T}}^2(\mathbf{z}_1, \mathbf{z}_2) = \psi(\mathbf{z}_1 \ominus \mathbf{z}_2)^t \cdot [\Sigma^\psi]^{-1} \cdot \psi(\mathbf{z}_1 \ominus \mathbf{z}_2),$$

is invariant under the choice of full-rank log-ratio representation $\psi(\cdot)$.

Proof:

To show this proposition, it suffices to observe that from Corollary 3 and the proof of Proposition 4 we have $\mathbf{H} \cdot \boldsymbol{\Sigma}^{c^-} \cdot \mathbf{H} = \mathbf{H} \cdot \boldsymbol{\Psi}^t \cdot \boldsymbol{\Sigma}^{\Psi^{-1}} \cdot \boldsymbol{\Psi} \cdot \mathbf{H} = \boldsymbol{\Psi}^t \cdot \boldsymbol{\Sigma}^{\Psi^{-1}} \cdot \boldsymbol{\Psi}$ so that

$$\begin{aligned} d_{M,\mathbf{T}}^2(\mathbf{z}_1, \mathbf{z}_2) &= \log(\mathbf{z}_1 \ominus \mathbf{z}_2)^t \cdot \mathbf{H}^t \cdot \boldsymbol{\Sigma}^{c^-} \cdot \mathbf{H} \cdot \log(\mathbf{z}_1 \ominus \mathbf{z}_2) \\ &= \log(\mathbf{z}_1 \ominus \mathbf{z}_2)^t \cdot \boldsymbol{\Psi}^t \cdot \boldsymbol{\Sigma}^{\Psi^{-1}} \cdot \boldsymbol{\Psi} \cdot \log(\mathbf{z}_1 \ominus \mathbf{z}_2) \\ &= -2 \log(\mathbf{z}_1 \ominus \mathbf{z}_2)^t \cdot \boldsymbol{\Psi}^t \cdot \boldsymbol{\Psi}^{-t} \cdot \hat{\mathbf{T}}^- \cdot \boldsymbol{\Psi}^- \cdot \boldsymbol{\Psi} \cdot \log(\mathbf{z}_1 \ominus \mathbf{z}_2) \end{aligned}$$

an expression which does not depend on the log-ratio representation at all. \square

Proposition 6 is a direct consequence of the invariance property of the Mahalanobis distance.

Proposition 6 (Invariance of the normal distribution) *The probability density function of the normal distribution on the simplex with center \mathbf{m} and variation matrix \mathbf{T} ,*

$$f_{\mathbf{z}}(\mathbf{z}) = (2\pi|\boldsymbol{\Sigma}^\psi|)^{(1-D)/2} \exp \left[-\frac{1}{2} d_{M,\mathbf{T}}^2(\mathbf{z}, \mathbf{m}) \right],$$

does not depend on the choice of full-rank log-ratio representation $\psi(\cdot)$.

Analogous results are available for the case when the log-ratio transform is not full-rank. In that case a the determinant $|\boldsymbol{\Sigma}^\psi|$ needs to be generalised to the product of its non-zero eigenvalues. This invariance (Mateu-Figueras et al., 2013) is a direct consequence of the preceding proposition 5 and the fact that the determinant of a matrix is one of its invariants.

A.2 Spatial results

Definition 3 (Compositional random function) *A vector-valued random function $\mathbf{Z} = [Z_1, Z_2, \dots, Z_D]$ on a spatial domain $\mathcal{D} \subset \mathbb{R}^p$, is called compositional if for each $x \in \mathcal{D}$ the vector of random variables $\mathbf{Z}(x) = [Z_1(x), Z_2(x), \dots, Z_D(x)]$ shows the relative importance of a set of parts forming a total of interest.*

Definition 4 (Regionalized composition) *Given a set of locations $\{x_1, x_2, \dots, x_N\}$, a regionalized data set $\{\mathbf{z}_1, \mathbf{z}_2, \dots, \mathbf{z}_N\}$ with $\mathbf{z}_i = \mathbf{z}(x_i) = [z_1(x_i), \dots, z_D(x_i)] = [z_{1i}, \dots, z_{Di}]$, $i = 1, 2, \dots, N$ is called a regionalized composition, if z_{ki} represents the relative importance of part k with respect to the set of components considered at location x_i .*

Proposition 7 (log-ratio representation of the spatial structure) *Let $\mathbf{Z} = [z_{kn}] = [z_k(x_n)]$, $k = 1, 2, \dots, D$, $n = 1, 2, \dots, N$, be a regionalized compositional data set with N locations x_n and D parts, and $\psi(\cdot)$ be a full-rank log-ratio transformation. Then, for each lag h , the variogram of the log-ratio representation can be obtained from the empirical variation-variogram $\hat{\mathbf{T}}(h)$ as $\hat{\Sigma}^\psi(h) = -\frac{1}{2}\Psi \cdot \hat{\mathbf{T}}(h) \cdot \Psi^t$, or from the clr-variogram matrix as $\hat{\Gamma}^\psi(h) = \Psi \cdot \hat{\Gamma}^c(h) \cdot \Psi^t$.*

This is a direct consequence of propositions 3 and 4.

Proposition 8 (Equivalence of the spatial structure) *Let $\mathbf{Z} = [z_{kn}] = [z_k(x_n)]$, $k = 1, 2, \dots, D$, $n = 1, 2, \dots, N$, be a regionalized compositional data set with N locations x_n and D parts, and $\psi_1(\cdot)$ and $\psi_2(\cdot)$ be two full-rank log-ratio transformations. Then, for each lag h , the empirical variograms $\hat{\Gamma}^{\psi_1}(h)$ and $\hat{\Gamma}^{\psi_2}(h)$ are related through the linear relationship*

$$\hat{\Gamma}^{\psi_2}(h) = \mathbf{A}_{12} \cdot \hat{\Gamma}^{\psi_1}(h) \cdot \mathbf{A}_{12}^t \quad (35)$$

with matrix $\mathbf{A}_{12} = \Psi_2 \cdot \Psi_1^{-1}$ square and invertible.

Proof: From proposition 7, it follows that $\hat{\Gamma}^{\psi_2}(h) = \Psi_2 \cdot \hat{\Gamma}^c(h) \cdot \Psi_2^t$; and because of Eq. (??), $\hat{\Gamma}^c(h) = \Psi_1^{-1} \cdot \hat{\Gamma}^{\psi_1}(h) \cdot \Psi_1^{-t}$. Therefore

$$\hat{\Gamma}^{\psi_2}(h) = \Psi_2 \cdot \Psi_1^{-1} \cdot \hat{\Gamma}^{\psi_1}(h) \cdot \Psi_1^{-t} \cdot \Psi_2^t,$$

which proves the desired equality because $\mathbf{A}_{12}^t = \Psi_1^{-t} \cdot \Psi_2^t$. \square

Since proposition 8 holds for all lags, it is normal to require that any fitted model satisfies the same relation. This is automatically satisfied if a linear model of coregionalization $\mathbf{T}(h|\boldsymbol{\theta})$ is fitted to the variation-variograms and then recast to each of the two log-ratio representations by using proposition 7.

Proposition 9 (Invariance of the cokriging predictor and errors) *Let $\mathbf{Z} = [z_{kn}] = [z_k(x_n)]$, $k = 1, 2, \dots, D$, $n = 1, 2, \dots, N$, be a regionalized compositional data set with N locations x_n and D parts, and $\psi_1(\cdot)$ and $\psi_2(\cdot)$ be two full-rank log-ratio transformations. Then, the corresponding cokriging predictors $\hat{\zeta}_1(x_0)$ and $\hat{\zeta}_2(x_0)$ of the log-ratio transformed composition $\zeta_i(x_0) = \psi_i(\mathbf{Z}(x_0))$ satisfy*

$$\hat{\zeta}_2(x_0) = \mathbf{A}_{12} \cdot \hat{\zeta}_1(x_0),$$

so that

$$\psi_1^{-1}(\hat{\zeta}_1(x_0)) = \psi_2^{-1}(\hat{\zeta}_2(x_0)) =: \hat{\mathbf{z}}(x_0)$$

gives a predicted composition independent of the log-ratio representation used in the computations. Moreover, the corresponding cokriging error covariance matrices \mathbf{S}_1 and \mathbf{S}_2 are related by

$$\mathbf{S}_2^K = \mathbf{A}_{12} \cdot \mathbf{S}_1^K \cdot \mathbf{A}_{12}^t$$

with $\mathbf{A}_{12} = \mathbf{\Psi}_2 \cdot \mathbf{\Psi}_1^-$, for all forms of cokriging (simple, ordinary, universal and cokriging with a trend) at all locations x_0 , if both are derived from the same linear model of coregionalization $\mathbf{T}(h|\boldsymbol{\theta})$.

Proof: We will first consider the case of simple cokriging (SK) under the assumption of second-order stationarity. In both log-ratio representations, the SK predictor is of the form

$$\hat{\zeta}(x_0) = \sum_{n=1}^N \lambda_n^t \zeta(x_n) = \boldsymbol{\Lambda}^t \mathbf{Z}, \quad (36)$$

where $\mathbf{Z} = [\zeta(x_1); \zeta(x_2); \dots; \zeta(x_N)]$ is the concatenated vector of all log-ratio transformed observations $\zeta(x_n) = \mathbf{\Psi} \log \mathbf{z}(x_n)$, and $\boldsymbol{\Lambda} = [\boldsymbol{\lambda}_1; \boldsymbol{\lambda}_2; \dots; \boldsymbol{\lambda}_N]$ is the block matrix of all cokriging weight matrices, which are obtained as Myers (1982),

$$\boldsymbol{\Lambda} = \underbrace{\begin{bmatrix} \boldsymbol{\Gamma}_{11} & \boldsymbol{\Gamma}_{12} & \cdots & \boldsymbol{\Gamma}_{1N} \\ \boldsymbol{\Gamma}_{21} & \boldsymbol{\Gamma}_{22} & \cdots & \boldsymbol{\Gamma}_{2N} \\ \vdots & \vdots & \ddots & \vdots \\ \boldsymbol{\Gamma}_{N1} & \boldsymbol{\Gamma}_{N2} & \cdots & \boldsymbol{\Gamma}_{NN} \end{bmatrix}}_{\mathbf{W}^{-1}}^{-1} \underbrace{\begin{bmatrix} \boldsymbol{\Gamma}_{10} \\ \boldsymbol{\Gamma}_{20} \\ \vdots \\ \boldsymbol{\Gamma}_{N0} \end{bmatrix}}_{\mathbf{W}_0} = \mathbf{W}^{-1} \mathbf{W}_0$$

where each block $\boldsymbol{\Gamma}_{nm} = \boldsymbol{\Gamma}(h|\boldsymbol{\theta}) = -\frac{1}{2} \mathbf{\Psi} \mathbf{T}(h|\boldsymbol{\theta}) \mathbf{\Psi}^t$ using the fitted model $\mathbf{T}(h|\boldsymbol{\theta})$. With the same notation, the SK error covariance is given by

$$\mathbf{S}^K = \boldsymbol{\Gamma}_{00} - \boldsymbol{\Lambda}^t \mathbf{W}_0 = \boldsymbol{\Gamma}_{00} - \mathbf{W}_0^t \mathbf{W}^{-1} \mathbf{W}_0.$$

If we now consider these matrices obtained with the two log-ratio representa-

tions, and taking Eq. (35) into account,

$$\begin{aligned}
\mathbf{W}^{(2)} &= \begin{bmatrix} \boldsymbol{\Gamma}_{11}^{(2)} & \boldsymbol{\Gamma}_{12}^{(2)} & \cdots & \boldsymbol{\Gamma}_{1N}^{(2)} \\ \boldsymbol{\Gamma}_{21}^{(2)} & \boldsymbol{\Gamma}_{22}^{(2)} & \cdots & \boldsymbol{\Gamma}_{2N}^{(2)} \\ \vdots & \vdots & \ddots & \vdots \\ \boldsymbol{\Gamma}_{N1}^{(2)} & \boldsymbol{\Gamma}_{N2}^{(2)} & \cdots & \boldsymbol{\Gamma}_{NN}^{(2)} \end{bmatrix} \\
&= \begin{bmatrix} \mathbf{A}_{12}\boldsymbol{\Gamma}_{11}^{(1)}\mathbf{A}_{12}^t & \mathbf{A}_{12}\boldsymbol{\Gamma}_{12}^{(1)}\mathbf{A}_{12}^t & \cdots & \mathbf{A}_{12}\boldsymbol{\Gamma}_{1N}^{(1)}\mathbf{A}_{12}^t \\ \mathbf{A}_{12}\boldsymbol{\Gamma}_{21}^{(1)}\mathbf{A}_{12}^t & \mathbf{A}_{12}\boldsymbol{\Gamma}_{22}^{(1)}\mathbf{A}_{12}^t & \cdots & \mathbf{A}_{12}\boldsymbol{\Gamma}_{2N}^{(1)}\mathbf{A}_{12}^t \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{A}_{12}\boldsymbol{\Gamma}_{N1}^{(1)}\mathbf{A}_{12}^t & \mathbf{A}_{12}\boldsymbol{\Gamma}_{N2}^{(1)}\mathbf{A}_{12}^t & \cdots & \mathbf{A}_{12}\boldsymbol{\Gamma}_{NN}^{(1)}\mathbf{A}_{12}^t \end{bmatrix} \\
&= \mathbf{A}\mathbf{W}^{(1)}\mathbf{A}^t,
\end{aligned} \tag{37}$$

where $\mathbf{A} = \text{diag}(\mathbf{A}_{12}, \mathbf{A}_{12}, \dots, \mathbf{A}_{12})$ and similiary,

$$\mathbf{W}_0^{(2)} = \mathbf{A}\mathbf{W}_0^{(1)}\mathbf{A}_{12}^t. \tag{38}$$

Now substituting (37) and (38) into the expression for the weights

$$\begin{aligned}
\boldsymbol{\Lambda}^{(2)} &= [\mathbf{W}^{(2)}]^{-1}\mathbf{W}_0^{(2)} = [\mathbf{A}\mathbf{W}^{(1)}\mathbf{A}^t]^{-1}\mathbf{A}\mathbf{W}_0^{(1)}\mathbf{A}_{12}^t = \\
&= \mathbf{A}^{-t}[\mathbf{W}^{(1)}]^{-1}\mathbf{A}^{-1}\mathbf{A}\mathbf{W}_0^{(1)}\mathbf{A}_{12}^t = \mathbf{A}^{-t}[\mathbf{W}^{(1)}]^{-1}\mathbf{W}_0^{(1)}\mathbf{A}_{12}^t = \\
&= \mathbf{A}^{-t}\boldsymbol{\Lambda}^{(1)}\mathbf{A}_{12}^t,
\end{aligned} \tag{39}$$

which, due to the block-diagonal structure of \mathbf{A} , implies that the cokriging weight matrices of each datum satisfy

$$\boldsymbol{\lambda}_n^{(2)} = \mathbf{A}_{12}^{-t}\boldsymbol{\lambda}_n^{(1)}\mathbf{A}_{12}^t$$

Finally substituting these weights into the SK predictor of the second log-ratio representation, and taking into account Eq. (34) between the data,

$$\begin{aligned}
\hat{\boldsymbol{\zeta}}_2(x_0) &= \sum_{n=1}^N [\boldsymbol{\lambda}_n^{(2)}]^t \boldsymbol{\zeta}_2(x_n) = \sum_{n=1}^N (\mathbf{A}_{12}^{-t}\boldsymbol{\lambda}_n^{(1)}\mathbf{A}_{12}^t)^t \mathbf{A}_{12}\boldsymbol{\zeta}_1 = \\
&= \sum_{n=1}^N \mathbf{A}_{12}[\boldsymbol{\lambda}_n^{(1)}]^t \mathbf{A}_{12}^{-1}\mathbf{A}_{12}\boldsymbol{\zeta}_1 = \mathbf{A}_{12} \sum_{n=1}^N [\boldsymbol{\lambda}_n^{(1)}]^t \boldsymbol{\zeta}_1 = \mathbf{A}_{12}\hat{\boldsymbol{\zeta}}_1(x_0),
\end{aligned}$$

thus establishing the identity between the cokriging predictors. To derive the relation for the cokriging error covariance, the same strategy can be used to express the error in terms of the second log-ratio representation as a function

of that in terms of the first representation,

$$\begin{aligned}
\mathbf{S}_{(2)}^K &= \mathbf{\Gamma}_{00}^{(2)} - [\mathbf{\Lambda}^{(2)}]^t \mathbf{W}_0^{(2)} = \mathbf{A}_{12} \mathbf{\Gamma}_{00}^{(1)} \mathbf{A}_{12}^t - [\mathbf{A}^{-t} \mathbf{\Lambda}^{(1)} \mathbf{A}_{12}^t]^t \mathbf{A} \mathbf{W}_0^{(1)} \mathbf{A}_{12}^t = \\
&= \mathbf{A}_{12} \mathbf{\Gamma}_{00}^{(1)} \mathbf{A}_{12}^t - \mathbf{A}_{12} [\mathbf{\Lambda}^{(1)}]^t \mathbf{A}^{-1} \mathbf{A} \mathbf{W}_0^{(1)} \mathbf{A}_{12}^t = \\
&= \mathbf{A}_{12} \mathbf{\Gamma}_{00}^{(1)} \mathbf{A}_{12}^t - \mathbf{A}_{12} [\mathbf{\Lambda}^{(1)}]^t \mathbf{W}_0^{(1)} \mathbf{A}_{12}^t = \mathbf{A}_{12} \left[\mathbf{\Gamma}_{00}^{(1)} - [\mathbf{\Lambda}^{(1)}]^t \mathbf{W}_0^{(1)} \right] \mathbf{A}_{12}^t = \\
&= \mathbf{A}_{12} \mathbf{S}_{(1)}^K \mathbf{A}_{12}^t,
\end{aligned}$$

which proves the desired equivalence.

For the remaining cases of cokriging (which will be grouped under the name of universal cokriging, UK), the log-ratio mean is assumed to have the form

$$\boldsymbol{\mu}(x) = \sum_{l=1}^L g_l(x) \mathbf{b}_l$$

with the typical cases $L = 1$ and $g_1(x) \equiv 1$ (for ordinary cokriging), $g_l(x) = x^{l-1}$ up to the desired order L (universal cokriging), or $L = 1$ and $g_1(x)$ an arbitrary function available everywhere in the estimation domain (for cokriging with a trend). In any case, the UK predictor has the same form (Eq. 36), where the weights are obtained by solving the system

$$\mathbf{W} \mathbf{\Lambda} = \mathbf{W}_0$$

subject to the L unbiasedness conditions

$$\sum_{n=1}^N g_l(x_n) \boldsymbol{\lambda}_n^t = g_l(x_0) \mathbf{I}_{D-1}, \quad l = 1, 2, \dots, L.$$

where \mathbf{I}_{D-1} is the identity matrix of size $(D-1)$, the dimension of the composition. It is known (Myers, 1982; Tolosana-Delgado, 2006) that this is equivalent to solving an extended system of equations

$$\mathbf{W}_e \mathbf{\Lambda}_e = \mathbf{W}_{e0} \tag{40}$$

where

$$\mathbf{W}_e = \begin{bmatrix} \mathbf{W} & \mathbf{G} \\ \mathbf{G}^t & 0_{\mathbf{I}_{L(D-1)}} \end{bmatrix}, \quad \mathbf{W}_{e0} = \begin{bmatrix} \mathbf{W}_0 \\ \mathbf{G}_0^t \end{bmatrix}, \quad \mathbf{\Lambda}_e = \begin{bmatrix} \mathbf{\Lambda} \\ \mathbf{N} \end{bmatrix},$$

with $\mathbf{N}^t = [\boldsymbol{\nu}_1; \boldsymbol{\nu}_2; \dots; \boldsymbol{\nu}_L]$ the Lagrange multipliers for each unbiasedness condition, and $\mathbf{G}^t = [\mathbf{G}_1^t; \mathbf{G}_2^t; \dots; \mathbf{G}_N^t]$ with

$$\mathbf{G}_i = [g_1(x_i) \mathbf{I}_{D-1}; g_2(x_i) \mathbf{I}_{D-1}; \dots; g_L(x_i) \mathbf{I}_{D-1}], \quad i = 0, 1, \dots, N.$$

The UK error covariance matrix is then shown to be

$$\mathbf{S}^K = \mathbf{\Gamma}_{00} - \mathbf{\Lambda}_e^t \mathbf{W}_{e0} = \mathbf{\Gamma}_{00} - \mathbf{W}_{e0}^t \mathbf{W}_e^{-1} \mathbf{W}_{e0}$$

Since the UK and SK system of equations, predictors and errors have analogous forms, the proposition for the case of UK can be proved by showing that, if the extended matrices satisfy Eqs. (37)-(39), then they satisfy the UK system of equations (Eq. 40) as well. That is, if $\mathbf{W}_e^{(2)} = \mathbf{A} \mathbf{W}_e^{(1)} \mathbf{A}^t$ (Eq. 37) and $\mathbf{\Lambda}_e^{(2)} = \mathbf{A}^{-t} \mathbf{\Lambda}_e^{(1)} \mathbf{A}_{12}^t$ (Eq. 39), then in Eq. (40), becomes

$$\mathbf{W}_e^{(2)} \mathbf{\Lambda}_e^{(2)} = [\mathbf{A} \mathbf{W}_e^{(1)} \mathbf{A}^t] [\mathbf{A}^{-t} \mathbf{\Lambda}_e^{(1)} \mathbf{A}_{12}^t] = \mathbf{A} \mathbf{W}_e^{(1)} \mathbf{\Lambda}_e^{(1)} \mathbf{A}_{12}^t = \mathbf{A} \mathbf{W}_{e0}^{(1)} \mathbf{A}_{12}^t = \mathbf{W}_{e0}^{(2)},$$

which holds given Eq. (38). \square

For an integral

$$I = \int_{\mathbb{R}^{D-1}} g(\mathbf{Z}) \phi(\mathbf{Z} | \hat{\boldsymbol{\zeta}}, \mathbf{S}^K) d\mathbf{Z},$$

where $\hat{\boldsymbol{\zeta}}$ is a cokriging predictor with cokriging error covariance \mathbf{S}^K , a multivariate Gauss-Hermite quadrature of order k with weights w_1, w_2, \dots, w_k and quadrature points u_1, u_2, \dots, u_k , can be used to obtain the approximation

$$\hat{I} = \sum_{i_1=1}^k \sum_{i_2=1}^k \cdots \sum_{i_{D-1}=1}^k w_{i_1} w_{i_2} \cdots w_{i_{D-1}} g(\boldsymbol{\zeta}(i_1, i_2, \dots, i_{D-1}))$$

with quadrature vectors

$$\boldsymbol{\zeta}(i_1, i_2, \dots, u_{D-1}) =: \hat{\boldsymbol{\zeta}} + \sqrt{2} \cdot \mathbf{R} \cdot \mathbf{u}_{[i_1, i_2, \dots, i_{D-1}]}$$

where \mathbf{R} is any matrix satisfying $\mathbf{R} \cdot \mathbf{R}^t = \mathbf{S}^K$ and $\mathbf{u}_{[i_1, i_2, \dots, i_{D-1}]} = [u_{i_1}, u_{i_2}, \dots, u_{i_{D-1}}]$,

We lastly establish the relationship between quadratures for distinct log-ratio representations $\psi_1(\cdot)$ and $\psi_2(\cdot)$. The weights w_1, w_2, \dots, w_k and quadrature points u_1, u_2, \dots, u_k do not depend on the choice of log-ratio representation. If $\hat{\boldsymbol{\zeta}}_i$ is the predictor using the i -th log-ratio representation, then by Proposition 9 the representations $\hat{\boldsymbol{\zeta}}_1$ and $\hat{\boldsymbol{\zeta}}_2$ are related by $\mathbf{A}_{12} \cdot \hat{\boldsymbol{\zeta}}_1 = \hat{\boldsymbol{\zeta}}_2$.

The spectral decomposition of the cokriging error covariance matrix \mathbf{S}_1^K is given by $\mathbf{S}_1^K = \mathbf{V}_1 \cdot \mathbf{D}_1 \cdot \mathbf{V}_1^t$, where \mathbf{D}_1 is a diagonal matrix and \mathbf{V}_1 is an orthogonal matrix of eigenvectors then $\mathbf{R}_1 = \mathbf{V}_1 \cdot \mathbf{D}_1^{1/2} \cdot \mathbf{V}_1^t$ is a square root of \mathbf{S}_1^K and so from the congruence we have

$$\mathbf{S}_2^K = \mathbf{A}_{12} \cdot \mathbf{S}_1^K \cdot \mathbf{A}_{12}^t = \mathbf{A}_{12} \cdot (\mathbf{V}_1 \cdot \mathbf{D}_1 \cdot \mathbf{V}_1^t) \cdot \mathbf{A}_{12}^t.$$

This expression can be rewritten as

$$\mathbf{S}_2^K = \mathbf{A}_{12} \cdot \mathbf{V}_1 \cdot \mathbf{D}_1^{1/2} \cdot \mathbf{V}_1^t \cdot (\mathbf{A}_{12}^{-1} \cdot \mathbf{A}_{12}) \cdot \mathbf{V}_1 \cdot \mathbf{D}_1^{1/2} \cdot \mathbf{V}_1^t \cdot \mathbf{A}_{12}^t = (\mathbf{A}_{12} \cdot \mathbf{R}_1 \cdot \mathbf{A}_{12}^{-1}) \cdot \mathbf{A}_{12} \cdot \mathbf{R}_1 \cdot \mathbf{A}_{12}^t$$

and so

$$\mathbf{R}_2 = \mathbf{A}_{12} \cdot \mathbf{R}_1 \cdot \mathbf{A}_{12}^t$$

is a square root of \mathbf{S}_2 if and only if $\mathbf{A}_{12}^t = \mathbf{A}_{12}^{-1}$, that is \mathbf{A}_{12} is an orthogonal matrix. In that case the quadrature vectors $\zeta(i_1, i_2, \dots, i_{D-1})$ are related by

$$\zeta_{(2)}(i_1, i_2, \dots, i_{D-1}) = \zeta_2 + \mathbf{R}_2 \cdot \mathbf{u}_{[i_1, i_2, \dots, i_{D-1}]} \quad (41)$$

$$= \mathbf{A}_{12} \cdot \zeta_1 + \mathbf{A}_{12} \cdot \mathbf{R}_1 \cdot \mathbf{A}_{12}^t \cdot \mathbf{u}_{[i_1, i_2, \dots, i_{D-1}]} \quad (42)$$

$$= \mathbf{A}_{12} \cdot (\zeta_1 + \mathbf{R}_1 \cdot \mathbf{v}_{[i_1, i_2, \dots, i_{D-1}]}). \quad (43)$$

where $\mathbf{v}_{[i_1, i_2, \dots, i_{D-1}]} = \mathbf{A}_{12} \cdot \mathbf{u}_{[i_1, i_2, \dots, i_{D-1}]}$.

B Appendix: Compositional Geostatistics Workflow

B.1 Interpolation

1. perform both classical and compositional exploratory analysis (section 3.3)
2. compute variation-variograms of the regionalized composition (Eq. 24)
3. fit a valid model (section 5.2); models such as the linear model of coregionalization or the minimum/maximum autocorrelation factors are useful
4. recast both the experimental and the model variation-variograms via other log-ratio transform with respectively Eqs. (25) and (27), in order to assess that the model fits the data reasonably well in these other reference systems
5. choose one of these alternative log-ratio transforms, and compute the scores of the data (Eq. 13)
6. apply cokriging to the log-ratio scores with variogram model expressed in the same log-ratios; store cokriging covariance error matrices if cross-validation or Gauss-Hermite quadratures is desired
7. backtransform the predicted values
8. if unbiased estimates of the mass of each component are required, estimate them through Gauss-Hermite quadratures (Eq. 30) or follow the procedure in the next section

9. further products (maps, cross-validation, block models, etc) can be derived from individual components of the composition or from relevant logratios; cross-validation studies should focus on multivariate quantities and pairwise logratio plots (section 6.2).

Steps (2) and (3) can alternatively be applied to a particular log-ratio transformed data. In this case, step (4) should also explore the fit of the model to the variation-variograms, and step (5) can be applied to the same logratio set as in step (2). This is the strategy followed in the paper, where all calculations were primarily done with the alr transformed data.

B.2 Simulation

1. apply a log-ratio transformation to the data, then transform the scores via multivariate Gaussian anamorphosis, such as the flow anamorphosis (section 7.2)
2. estimate direct and cross-variograms of the gaussian scores
3. fit a valid joint model to these variograms
4. apply conditional simulation algorithms to produce simulations of the gaussian scores
5. transform the simulated gaussian scores to logratio scores with the inverse Gaussian anamorphosis, then backtransform the logratio scores to compositions
6. post-process simulations as desired, e.g. produce point-wise estimates of non-linear quantities (Eq 29), upscale them to block averages (Eqs. 31-33) or produce maps.

References

- Aitchison, J. (1982). The statistical analysis of compositional data (with discussion). *Journal of the Royal Statistical Society, Series B (Statistical Methodology)* 44(2), 139–177.
- Aitchison, J. (1986). *The Statistical Analysis of Compositional Data*. Monographs on Statistics and Applied Probability. Chapman & Hall Ltd., London (UK). (Reprinted in 2003 with additional material by The Blackburn Press). 416 p.

- Angerer, T. and Hagemann, S. (2010). The BIF-hosted high-grade iron ore deposits in the Archean Koolyanobbing Greenstone Belt, Western Australia: Structural control on synorogenic- and weathering-related magnetite-, hematite- and goethite-rich iron ore. *Economic Geology* 105(3), 917–945.
- Barnett, R. M., Manchuk, J. G. and Deutsch, C.V. (2014). Projection pursuit multivariate transform. *Mathematical Geosciences* 46 (2), 337–360.
- Bivand, R.S., E. Pebesma and V. Gomez-Rubio (2013). *Applied Spatial Data Analysis with R*. Springer, New York, NY (USA). 405 p.
- Barceló-Vidal, C. (2003). When a data set can be considered compositional?. In S. Thi'o-Henestrosa and J.A. Martin-Fern'andez (Eds.), *Proceedings of CoDaWork'03, The 1st Compositional Data Analysis Workshop*. Universitat de Girona.
- Chayes, F. (1960). On correlation between variables of constant sum. *Journal of Geophysical Research* 65(12), 4185–4193.
- Chilés J. P. and Delfiner P. (1999). *Geostatistics..* John Wiley and Sons, New York, NY (USA). 695 p.
- Cressie, N. (1991). *Statistics for Spatial Data*. John Wiley and Sons, New York, NY (USA). 900 p.
- Egozcue, J.J., V. Pawlowsky-Glahn, G. Mateu-Figueras and C. Barceló-Vidal (2008). Isometric logratio transformations for compositional data analysis. *Mathematical Geosciences* 35(3), 279–300.
- Filzmoser, P. and K. Hron (2008). Outlier detection for compositional data using robust methods. *Mathematical Geosciences* 40(3), 233–248.
- Gelfand, A. E. and S. Banerjee (2010). *Multivariate Spatial Process Models*, Chapter 28, pp. 495–515.
- Geovariances (2017). *Isatis Geostatistical Software*. Avon, France.
- Griffin, A. C. (1981). *Structure and Iron Ore deposition in the Archaean Koolyanobbing Greenstone belt, Western Australia*. Geological Society of Australia, special publication 7, pp. 429–438.
- Mueller, U. A. and E. C. . Grunsky (2016). Multivariate spatial analysis of lake sediment geochemical data; Melville Peninsula, Nunavut, Canada. *Applied Geochemistry* 75, 247–262.

- Journal, A. G. and C. J. Huijbregts (1978). *Mining Geostatistics*. Academic Press, London (UK). 600 p.
- Lark, R. M. and T. F. A. Bishop (2007). Cokriging particle size fractions of the soil. *European Journal of Soil Science* 58(3), 763–774.
- Leuangthong, O. and Deutsch, C.V. (2003). Stepwise conditional transformation for simulation of multiple variables. *Mathematical Geology* 35 (2), 155–173.
- Mateu-Figueras, G., V. Pawlowsky-Glahn, and J. J. Egozcue (2013). The normal distribution in some constrained sample spaces. *SORT - Statistics and Operations Research Transactions* 37(1), 29–56.
- Matheron, G. (1963). Principles of geostatistics. *Economic Geology* 58, 1246–1266.
- Matheron, G. (1965). *Les variables régionalisées et leur estimation—une application de la théorie des fonctions aléatoires aux sciences de la nature*. Masson et Cie., Paris (F). 305 p.
- Matheron, G. (1971). The theory of regionalized variables and its applications. Technical Report C-5, École Nationale Supérieure des Mines de Paris, Centre de Geostatistique et de Morphologie Mathématique, Fontainebleau (F). 211 p.
- Morales Boezio, M. N. (2010). *Estudo das metodologias alternativas da geostatística multivariada aplicadas a estimativa de teores de depósitos de ferro*. Ph. D. thesis, Universidade Federal do Rio Grande do Sul. Escola de Engenharia. Programa de Pós-Graduação em Engenharia Minas, Metalúrgica e de Materiais.
- Morales Boezio, M. N., J. F. Costa, and J. C. Koppe (2012, 09). Cokrigagem de razões logarítmicas aditivas (alr) na estimativa de teores em depósitos de ferro (Cokriging of additive log-ratios (alr) for grade estimation in iron ore deposits. *Rem: Revista Escola de Minas* 65, 401 – 412.
- Mueller, U., R. Tolosana-Delgado, and K. G. van den Boogaart (2014, October). Simulation of compositional data: A nickel-laterite case study. In R. Dimitrakopoulos (Ed.), *Advances in Orebody Modelling and Strategic Mine Planning*. AusIMM.
- Myers, D. E. (1982). Matrix formulation of co-kriging. *Mathematical Geology* 14(3), 249–257.

- Pawlowsky, V. (1984). On spurious spatial covariance between variables of constant sum. *Science de la Terre, Sér. Informatique* 21, 107–113.
- Pawlowsky, V. (1986). *Räumliche Strukturanalyse und Schätzung ortsabhängiger Kompositionen mit Anwendungsbeispielen aus der Geologie*. Ph. D. thesis, Fachbereich Geowissenschaften, Freie Universität Berlin, Berlin (D). 170 p.
- Pawlowsky, V. (1989). Cokriging of regionalized compositions. *Mathematical Geology* 21(5), 513–521.
- Pawlowsky-Glahn, V. and H. Burger (1992). Spatial structure analysis of regionalized compositions. *Mathematical Geology* 24(6), 675–691.
- Pawlowsky-Glahn, V. and J.J. Egozcue (2001). Geometric approach to statistical analysis on the simplex. *Stochastic Environmental Research and Risk Assessment* 15(5), 384–398.
- Pawlowsky-Glahn, V. and J.J. Egozcue (2002). BLU estimators and compositional data. *Mathematical Geology* 34(3), 259–274.
- Pawlowsky-Glahn, V. (2003). Statistical modelling on coordinates. In S. Thi'ohenestrosa and J.A. Martin-Fern'andez (Eds.), *Proceedings of CoDaWork'03, The 1st Compositional Data Analysis Workshop*. Universitat de Girona.
- Pawlowsky-Glahn, V. and R. A. Olea (2004). *Geostatistical Analysis of Compositional Data*. Number 7 in Studies in Mathematical Geology. Oxford University Press. Editor: DeGraffenreid, Jo Anne.
- Pawlowsky-Glahn, V., R. A. Olea, and J. C. Davis (1995). Estimation of regionalized compositions: A comparison of three methods. *Mathematical Geology* 27(1), 105–127.
- Rossi, M. E. and C. Deutsch (Eds.) (2014). *Mineral Resource Estimation*. Handbooks of Modern Statistical Methods. Springer.
- Sun, X. L., Y. J. Wu, H. L. Wang, Y. G. Zhao, and G. L. Zhang (2014). Mapping soil particle size fractions using compositional kriging, cokriging and additive log-ratio cokriging in two case studies. *Mathematical Geosciences* 46(4), 429–443.
- Tolosana-Delgado, R. (2006). *Geostatistics for constrained variables: positive data, compositions and probabilities. Application to environmental hazard monitoring*. Ph. D. thesis, Universitat de Girona (Spain). 198 p.

- Tolosana-Delgado, R. and K. G. v. d. Boogaart (2013). Joint consistent mapping of high-dimensional geochemical surveys. *Mathematical Geosciences* 45(8), 983–1004.
- Tolosana-Delgado, R., K. G. v. d. Boogaart, and V. Pawlowsky-Glahn (2009, June). Estimating and modeling variograms of compositional data with occasional missing variables in R. In *StatGIS'09, Geoinformatics for environmental surveillance Workshop*, Milos (Greece).
- Tolosana-Delgado, R., K. G. v. d. Boogaart, and V. Pawlowsky-Glahn (2011). Geostatistics for compositions. In V. Pawlowsky-Glahn and A. Buccianti (Eds.), *Compositional Data Analysis: Theory and Applications*, pp. 73–86. John Wiley & Sons. 378 p.
- Tolosana-Delgado, R., J. J. Egozcue, and V. Pawlowsky-Glahn (2008). Cokriging Of Compositions: logRatios And Unbiasedness. In J. M. Ortiz and X. Emery (Eds.), *Geostatistics Chile 2008*, pp. 299–308. Gecamin Ltd., Santiago, Chile, 2 vols, 1188 p.
- Tolosana-Delgado, R., U. Mueller, K. G. van den Boogaart, and C. Ward (2013). Block cokriging of a whole composition. pp. 267–277.
- Tolosana-Delgado, R., N. Otero, and V. Pawlowsky-Glahn (2005). Some basic concepts of compositional geometry. *Mathematical Geology* 37(7), 673–680.
- Tolosana-Delgado, R., U. Mueller, K. G. van den Boogaart, C. Ward and J. Gutzmer (2015). Improving processing by adaption to conditional geostatistical simulation of block compositions. *The Journal of The Southern African Institute of Mining and Metallurgy* 115(1), 13–26.
- van den Boogaart, K. G. R. Tolosana-Delgado and M. Bren (2009). compositions: Compositional Data Analysis package R package version 1.40-1.
- van den Boogaart, K. G. and R. Tolosana-Delgado, (2013). *Analysing Compositional Data with R*. Springer, Heidelberg (Germany). 280 p.
- van den Boogaart, K. G., R. Tolosana-Delgado, and U. Mueller (2017). An affine equivariant multivariate normal score transform for compositional data. *Mathematical Geosciences* 49(2), 231–252.
- Wackernagel, H. (2003). *Multivariate Geostatistics: An Introduction with Applications*. Springer, Berlin (Germany). 291 p.

- Walwoort, D. J. and J. J. de Gruijter (2001). Compositional Kriging: a Spatial Interpolation Method for Compositional Data. *Mathematical Geology* 33(8), 951–966.
- Ward, C. and U. Mueller (2012). *Multivariate Estimation Using Log Ratios: A Worked Alternative*, pp. 333–343. Quantitative Geology and Geostatistics. Springer.
- Ward, C. and U. Mueller (2013). Compositions, log ratios and bias - from grade control to resource. In *Iron Ore 2013 Shifting the paradigm*, pp. 313–320. The Australasian Institute of Mining and Metallurgy.